

WEAK MARKOV PROCESSES AS LINEAR SYSTEMS

ROLF GOHM


ABSTRACT. A noncommutative Fornasini-Marchesini system (a multi-variable version of a linear system) can be realized within a weak Markov process (a model for quantum evolution). For a discrete time parameter this is worked out systematically as a theory of representations of structure maps of a system by a weak process. We introduce subprocesses and quotient processes which can be described naturally by a suitable category of weak processes. A corresponding notion of cascade for processes induces a represented cascade of systems. We study the control theoretic notion of observability which turns out to be particularly interesting in connection with a cascade structure. As an application we gain new insights into stationary Markov chains where observability for the system is closely related to asymptotic completeness in the scattering theory of the chain. This motivates a general definition of asymptotic completeness in the category of weak processes.

1. INTRODUCTION

Recently there has been much attention for a certain multi-variable version of linear system theory which presents evolution equations of the form

$$\begin{aligned} x(\alpha k) &= A_k x(\alpha) + B_k u(\alpha), \\ y(\alpha) &= C x(\alpha) + D u(\alpha). \end{aligned}$$

This is called a noncommutative Fornasini-Marchesini system in [3, 4, 5]. Here $\alpha \in F_d^+$ where F_d^+ is the free semigroup with d generators (which we denote $1, \dots, d$ and $d \in \mathbb{N}$ or $d = \infty$). The elements in F_d^+ are words in the letters $1, \dots, d$, including the empty word 0. Composition is defined by concatenation of words, for example αk in the formula above is the concatenation of a word α and a generator $k \in \{1, \dots, d\}$. Now u, x, y are functions on F_d^+ taking values in vector spaces $\mathcal{U}, \mathcal{X}, \mathcal{Y}$ and $A_k : \mathcal{X} \rightarrow \mathcal{X}$, $B_k : \mathcal{U} \rightarrow \mathcal{X}$, $C : \mathcal{X} \rightarrow \mathcal{Y}$, $D : \mathcal{U} \rightarrow \mathcal{Y}$ are linear operators. For $d = 1$ we have $F_d^+ \simeq \mathbb{N}_0$ and this is the classical setting of linear system theory (state space models). The functions u, x, y are interpreted as input, internal state, output of the system.



It has been established that many important mathematical concepts and results in linear system theory generalize nicely for all d , see [2]. We mention two such concepts which will later be studied in this paper. The first is the *observability map*

$$\mathcal{O}_{C,A} := (C A^\alpha)_{\alpha \in F_d^+}$$

which is a linear map from $\mathcal{X}$ into the $\mathcal{Y}$ -valued functions on F_d^+ . (In this paper we use the convention that for any variables $X_1, \dots, X_d$ and a word $\alpha = \alpha_1 \dots \alpha_n \in F_d^+$ we have $X_\alpha := X_{\alpha_1} \dots X_{\alpha_n}$,

Date: March 6, 2013.

2000 Mathematics Subject Classification. Primary 46L53; Secondary 47A20, 93B07.

Key words and phrases. multi-variable system, weak Markov process, cascade, observability, asymptotic completeness.

$X^\alpha := X_{\alpha_n} \dots X_{\alpha_1}$, if $\alpha = 0$ we interpret it as an identity.) By studying the observability map we can find out what we are able to know about the internal space $\mathcal{X}$ by observing the output.

The second concept we want to mention here is the *transfer function* which is a description of how inputs are transferred into outputs. In the multi-variable context above the transfer function can be defined as a formal power series

$$\mathcal{T}(z) := \mathcal{T}(z_1, \dots, z_d) := \sum_{\alpha \in F_d^+} \mathcal{T}^\alpha z^\alpha := D + C \sum_{r=1}^{\infty} (zA)^{r-1} zB = D + C(I_{\mathcal{X}} - zA)^{-1} zB$$

where $z = (z_1 I_{\mathcal{X}}, \dots, z_d I_{\mathcal{X}})$ with indeterminates $z_1, \dots, z_d$ freely noncommuting among each other but commuting with the linear maps, $A = (A_1, \dots, A_d)^t$, $B = (B_1, \dots, B_d)^t$, and t denoting a transpose of the row vectors so A and B are column vectors of linear maps. In the classical case $d = 1$ we call the single variable z and then this reduces to a familiar formula which gives the transfer function as an analytic function. Explicitly

$$\mathcal{T}^\alpha = \begin{cases} D & \text{if } \alpha = 0 \\ C B_\alpha & \text{if } |\alpha| = 1 \\ C A_{\alpha_r} \dots A_{\alpha_2} B_{\alpha_1} & \text{if } \alpha = \alpha_1 \dots \alpha_r, r = |\alpha| \geq 2 \end{cases}$$

We come back to this at the end of Section 3. See also [2, 25] for further discussions of similar formulas.

Mathematical system theory is an abstraction from the physical dynamics. But of course its relevance to the real world depends on the fact that such a physical dynamics exists in the background. The basic idea behind our work is the proposal that the physical dynamics which implements a multi-variable system theory (such as the one sketched above) is described by processes describing quantum physics. In [24] we started a project to investigate such connections, compare further [17] for a recent generalization of the model in [24] and also [25] for some complementary work. In these papers the quantum processes are based on infinite tensor products which are natural from the point of view of approximating the Fock spaces of continuous time processes but which sometimes obscure the more fundamental level where the connections with operator and system theory originate. In the present paper we go directly to this level by investigating the proposal that it is the concept of ‘weak Markov processes’, worked out by Bhat and Parthasarathy in [8, 9] to catch the most basic features of quantum Markov processes, which provides a systematic and very general point of departure for our project. The connections to operator theory are evident here because the concept of a weak Markov process may be seen as operator theoretic dilation theory studied from a probabilistic point of view. Dilation theory will be mentioned only in side remarks in this paper however, the emphasis lies on the structure of the full process. The benefits of such studies go in both directions: access to operator and system theory tools for concrete quantum models and guidance for the development of the general theory from the questions arising in the models. To achieve such an interaction between the disciplines a more systematic development of the theory is needed. The following sketch of the contents of this paper should be read with this motivation in mind.

In Section 2 we start with a self-contained but concise description of the basic theory of weak Markov processes in discrete time: The dynamics is described by a row isometry and then from a co-invariant subspace certain structures emerge which can be given a quantum probabilistic interpretation, such as transition operators and weak filtrations. Most of this is well known but note how in the end we give an operational meaning to the elements of the free semigroup F_d^+ by interpreting them as certain measurement protocols.

In Section 3 we define the notion of a representation of structure maps A, B, C, D (as described above) by a weak Markov process and in this way we get an explicit systematic procedure to identify multi-variable linear systems (as described above) within quantum physical models. At this point it remains quite abstract. As a preparation for seeing representations of structure maps in use we first introduce in Section 4 the theory of subprocesses and quotient processes of (discrete weak Markov) processes and then show that within a suitably defined category of processes this can be reformulated as a short exact sequence. The main result here is a classification of extensions appearing in such short exact sequences by a construction which we call a γ -cascade of processes. There is a set of contractions

from which γ can be chosen which gives a parametrization of all the ways in which two processes can be put together as subprocess and quotient process, with $\gamma = 0$ yielding the direct sum.

Cascades are a well known concept for systems and their structure maps. The terminology of a γ -cascade for processes is justified in Section 5 where we show that within a γ -cascade of processes we have a representation of a cascade of the original systems and that in fact this construction is a natural tool to represent cascades of systems and of structure maps. Let us remark here that the notion of cascades and of more general networks of processes has been around for some time in a continuous time setting and this theory has been investigated intensely because of promising applications in quantum filtering and quantum control [26, 27]. There is no explicit comparison with continuous time systems in this paper, however we provide the basis for such a comparison by describing weak Markov processes in terms of product systems which suggests how to build the theory starting from continuous product systems. Our justification for concentrating on the discrete time setting here is the same as in Helton's seminal paper [29] in which he connects classical system theory ($d = 1$) with operator models and scattering theory, saying: 'We concentrate on discrete time systems because it is for these that the relationship is most clear' ([29], p.15).

In fact, very similarly to the path followed by Helton in [29] it is quite natural in our setting to connect observability of represented multi-variable linear systems with scattering in quantum physical models. We investigate observability in Section 6 and show that in a quantum model it can be given the same interpretation as in classical system theory, namely that by observing the outputs we have an indirect way to measure and to investigate those parts of the internal space which have the character of a black box. We show that we get a particularly interesting situation if we have a γ -cascade and if we use the input space of the subprocess as the output space of the cascade, so that looking for observability amounts to the question how much we can find out about the cascade by observations which only involve the subprocess. The extreme case when we can find out everything about the cascade by such special observations we call asymptotically complete.

We give a convenient criterion for asymptotic completeness in terms of asymptotic properties of the transition operator of the process which is a completely positive map. This criterion is a generalization of a criterion for asymptotic completeness [21, 22, 20] in a scattering theory for operator algebraic stationary quantum Markov processes introduced by Kümmerer and Maassen in [30]. It turns out that our definition of asymptotic completeness for short exact sequences of weak processes can be seen as a generalization of the definition of Kümmerer and Maassen. Such a connection has been hinted at in [24] but the theory developed in this paper allows us to explain the relevant facts in a much more conceptual and elegant way. The details are in Section 7. We include the general fact that normal invariant states lead to weak subprocesses and we work out how for a stationary Markov chain in an operator algebraic setting we can construct an associated weak process together with a subprocess (with one-dimensional internal space) and hence we can establish the connection to our previous results. This improved conceptual understanding of asymptotic completeness is an excellent example of the cross-fertilization between operator and system theory on the one hand and quantum probability on the other hand which we have in mind. It uses only a very special case of our general theory and suggests further work.

2. WEAK MARKOV PROCESSES

Unitary dynamics on a Hilbert space is the most basic way of describing quantum mechanical evolution. If causality is taken into account and one restricts the attention to observables belonging only to the future (or only to the past) then it becomes natural to study $*$ -endomorphisms of $\mathcal{B}(\mathcal{H})$, the algebra of bounded linear operators on a Hilbert space $\mathcal{H}$. A convincing argument in this direction is presented in [1], Section 1.2. From now on let $\theta : \mathcal{B}(\mathcal{H}) \rightarrow \mathcal{B}(\mathcal{H})$ be a $*$ -endomorphism and $\mathcal{H}$ a separable Hilbert space. Then it follows from the representation theory of $\mathcal{B}(\mathcal{H})$ that there exists a separable Hilbert space $\mathcal{P}$ and an isometry $V : \mathcal{H} \otimes \mathcal{P} \rightarrow \mathcal{H}$ such that for all $X \in \mathcal{B}(\mathcal{H})$

$$\theta(X) = V X \otimes \mathbb{1}_{\mathcal{P}} V^* .$$

We assume $\theta \neq 0$, then θ is automatically injective. See [32] for more details. If $(\epsilon_k)_{k=1}^d$ (with $d \in \mathbb{N}$ or $d = \infty$) is an orthonormal basis of $\mathcal{P}$ then with

$$V_k := V|_{\mathcal{H} \otimes \epsilon_k} \in \mathcal{B}(\mathcal{H}), \quad k = 1, \dots, d,$$

we can also write

$$\theta(X) = \sum_{k=1}^d V_k X V_k^*$$

(limits to be understood in the strong operator topology if $d = \infty$) which is called a *Kraus decomposition*. The V_k are isometries with orthogonal ranges, so alternatively (and with the same notation) we also think of V as a *row isometry*

$$V = (V_1, V_2, \dots, V_d) : \bigoplus_1^d \mathcal{H} \rightarrow \mathcal{H}.$$

V is called a *row unitary* if $\sum_{k=1}^d V_k V_k^* = \mathbb{1}$ and this is equivalent to θ being unital, i.e., $\theta(\mathbb{1}) = \mathbb{1}$. If this additional assumption is used in the following then it will always be explicitly stated.

Using induction we define row isometries $V^{(n)} : \mathcal{H} \otimes \bigotimes_1^n \mathcal{P} \rightarrow \mathcal{H}$ (where $n \in \mathbb{N}$) by $V^{(1)} := V$ and, for $\tilde{\xi} \in \mathcal{H} \otimes \bigotimes_1^{n-1} \mathcal{P}$ and $\eta \in \mathcal{P}$

$$V^{(n)}(\tilde{\xi} \otimes \eta) = V(V^{(n-1)}\tilde{\xi} \otimes \eta).$$

Later we need the following equivalent description for the adjoints: If $\xi \in \mathcal{H}$ and $V^*\xi = \sum_k \xi_k \otimes \eta_k \in \mathcal{H} \otimes \mathcal{P}$ then

$$V^{(n)*}\xi = \sum_k V^{(n-1)*}(\xi_k) \otimes \eta_k \in (\mathcal{H} \otimes \bigotimes_1^{n-1} \mathcal{P}) \otimes \mathcal{P} = \mathcal{H} \otimes \bigotimes_1^n \mathcal{P}.$$

It is not difficult to check that this implements the n -th power of θ :

$$\theta^n(X) = V^{(n)} X \otimes \mathbb{1}_{\bigotimes_1^n \mathcal{P}} V^{(n)*} = \sum_{\alpha \in F_d^+, |\alpha|=n} V_\alpha X V_\alpha^*.$$

Here the notation is $V_\alpha^* := (V_\alpha)^* = V_{\alpha_n}^* \dots V_{\alpha_1}^*$ if $\alpha = \alpha_1 \dots \alpha_n$ and $|\alpha| = n$ is the length of the word. This Kraus decomposition where a sum over all words of length n occurs gives the first connection to the multi-variable formalism sketched in Section 1.

Remark: The tensor products $(\bigotimes_1^n \mathcal{P})_{n \in \mathbb{N}}$ appearing in these formulas represent the (discrete) product system associated to the endomorphism θ and though we do not go into continuous time systems in this paper it is worth noting that the natural starting point to translate our results to continuous time would be to consider continuous product systems and in this way make the connection with the theory exposed in [1, 11, 23].

To introduce processes which resemble Markov processes from probability theory we need to specify a subspace $\mathfrak{h} \subset \mathcal{H}$ (by which we always mean a closed subspace if not otherwise stated). Let us denote the orthogonal projection onto $\mathfrak{h}$ by $p = P_{\mathfrak{h}}$. (In this paper we use consistently the notation $P_{\mathcal{L}}$ for the orthogonal projection onto a subspace $\mathcal{L}$.) Given $\mathfrak{h} \subset \mathcal{H}$ we have (with $n \in \mathbb{N}_0$) a family of normal $*$ -homomorphisms

$$\begin{aligned} J^{(n)} : \mathcal{B}(\mathfrak{h}) &\rightarrow \mathcal{B}(\mathcal{H}) \\ x &\mapsto \theta^n(xp) \end{aligned}$$

and the compressions $Z_n : \mathcal{B}(\mathfrak{h}) \rightarrow \mathcal{B}(\mathfrak{h})$ defined by

$$p J^{(n)}(x) p =: Z_n(x) p \quad [= J^{(0)}(Z_n(x))]$$

which are contractive completely positive maps. For the processes to be defined below the $J^{(n)}$ play the role of *noncommutative random variables* and the Z_n are *transition operators*.

The subspace $\mathfrak{h}$ of $\mathcal{H}$ is called *invariant* if $V_k \mathfrak{h} \subset \mathfrak{h}$ for all $k = 1, \dots, d$ and *co-invariant* if $V_k^* \mathfrak{h} \subset \mathfrak{h}$ for all $k = 1, \dots, d$. The importance of co-invariant subspaces in this context has been observed by

many, see for example [14] for various related topics. We note some useful properties equivalent to co-invariance.

Lemma 2.1. The following are equivalent:

- (1) $\theta(p)p = \theta(\mathbb{1})p$
- (2) $\mathfrak{h} \perp V(\mathfrak{h}^\perp \otimes \mathcal{P})$
- (3) $\mathfrak{h}$ is co-invariant

Let us write Z for Z_1 and state the following modification of the previous lemma:

Lemma 2.2. The following are equivalent:

- (1') $p \leq \theta(p)$
- (2') $\mathfrak{h} \subset V(\mathfrak{h} \otimes \mathcal{P})$
- (3') $p \leq \theta(\mathbb{1})$ and $\mathfrak{h}$ is co-invariant
- (4') $Z(\mathbb{1}_{\mathfrak{h}}) = \mathbb{1}_{\mathfrak{h}}$

Below we prove Lemma 2.1. It is then easy to get Lemma 2.2 by checking that (1') is nothing but (1) together with $p \leq \theta(\mathbb{1})$ (for $\ell = 1, \dots, 3$). The equivalence of (1') and (4') is immediate because, by definition, $Z(\mathbb{1}_{\mathfrak{h}}) = p\theta(p)|_{\mathfrak{h}}$.

If V is a row unitary then all the properties in Lemmas 2.1 and 2.2 are equivalent; for example (3) and (3') are equivalent for a row unitary because in this case we have $\theta(\mathbb{1}) = \mathbb{1}$. This means that for a row unitary we can always use the simpler statements (1')-(4') when dealing with co-invariant subspaces.

Proof. Consider an orthogonal projection q onto $q\mathcal{H}$. Then $\theta(q) = V(q \otimes \mathbb{1}_{\mathcal{P}})V^*$ is the orthogonal projection onto $V(q\mathcal{H} \otimes \mathcal{P})$. Hence $\theta(\mathbb{1} - p)$ projects onto $V(\mathfrak{h}^\perp \otimes \mathcal{P})$. This gives (1) $\Leftrightarrow$ (2).

Applying V^* to (2) we obtain $V^*\mathfrak{h} \perp \mathfrak{h}^\perp \otimes \mathcal{P}$, hence $V^*\mathfrak{h} \subset \mathfrak{h} \otimes \mathcal{P}$ and $V_k^*\mathfrak{h} \subset \mathfrak{h}$ for all $k = 1, \dots, d$ which is (3). Conversely, from (3) we get

$$\theta(p)p = \sum_1^d V_k p V_k^* p = \sum_1^d V_k V_k^* p = \theta(\mathbb{1})p$$

which is (1). □

The following definition is consistent with the terminology used by Bhat and Parthasarathy in [8, 9] where also continuous time and versions with C^* -subalgebras of $\mathcal{B}(\mathfrak{h})$ are considered which allows the inclusion of classical Markov processes by restricting to commutative subalgebras. We only consider discrete time steps and focus on the algebra $\mathcal{B}(\mathfrak{h})$ of all bounded operators. This allows a lean formulation from which further (probabilistic) features of the processes can be pulled out afterwards.

Definition 2.3. A *discrete weak Markov process* is a triple $(\mathcal{H}, V, \mathfrak{h})$ where V is a row isometry on a Hilbert space $\mathcal{H}$ which contains the Hilbert space $\mathfrak{h}$ as a co-invariant subspace with respect to V . Further we assume that

$$\mathcal{H} := \overline{\text{span}}\{V_\alpha \mathfrak{h} : \alpha \in F_d^+\}.$$

If V is a row unitary then we call the process *unital*.

For simplicity we refer to discrete weak Markov processes as processes. Note that if V is originally defined on a larger Hilbert space which contains the Hilbert space $\mathfrak{h}$ as a co-invariant subspace with respect to V then we can always restrict to a space $\mathcal{H}$ satisfying the additional assumption which ensures that $\mathcal{H}$ is the smallest invariant subspace containing $\mathfrak{h}$. Because $\mathfrak{h}$ is co-invariant it follows that $\mathcal{H}$ is even reducing in the original larger Hilbert space.

Remark: From the point of view of dilation theory V is the minimal isometric dilation of its compression to $\mathfrak{h}$ and as such it is determined by it up to unitary equivalence, see [33].

Let us now investigate the following increasing sequence of subspaces which is called the *weak filtration* associated to the process:

$$q_k := \theta^k(p), \quad p_k := \sup(q_0, \dots, q_k), \quad \mathfrak{h}_k := p_k \mathcal{H}$$

(for $k \in \mathbb{N}_0$). Then $p = p_0 \leq p_1 \leq p_2 \leq \dots$ and $\mathfrak{h} = \mathfrak{h}_0 \subset \mathfrak{h}_1 \subset \mathfrak{h}_2 \subset \dots$

Note that if the process is unital then $q_k = p_k$ for all k and the arguments simplify. For unital processes we have $\mathfrak{h}_n = V^{(n)}(\mathfrak{h} \otimes \bigotimes_1^n \mathcal{P})$ and the inclusion $\mathfrak{h}_n \subset \mathfrak{h}_{n+1}$ is mapped by $V^{(n+1)*}$ to an inclusion $\mathfrak{h} \otimes \bigotimes_1^n \mathcal{P} \subset \mathfrak{h} \otimes \bigotimes_1^{n+1} \mathcal{P}$. Hence in this case we have an identification of $\mathcal{H}$ with the inductive limit of the sequence $(\mathfrak{h} \otimes \bigotimes_1^n \mathcal{P})$.

Proposition 2.4. Let $(\mathcal{H}, V, \mathfrak{h})$ be a process. Then

- (a) $V(\mathfrak{h}_m \otimes \mathcal{P}) \subset \mathfrak{h}_{m+1}$ (for all $m \in \mathbb{N}_0$)
- (b) $Z_n = Z^n$ (for all $n \in \mathbb{N}$)
- (c) If $0 \leq m \leq n$ then $p_m V^{(n)} = q_m V^{(n)}$.
- (d) If $0 \leq m \leq n$ then

$$p_m J^{(n)}(x) p_m = J^{(m)}(Z^{n-m}(x))$$

(a) states an adaptedness property for the process and the filtration while (b), (c) and (d) resemble properties of Markov processes in classical probability and hence motivate the terminology ‘weak Markov process’. Versions of (d) appear in Bhat’s papers as the ‘weak Markov property’.

Proof. $\theta(p_m)$ is the projection onto $V(\mathfrak{h}_m \otimes \mathcal{P})$. Hence (a) is nothing but a reformulation of the obvious $\theta(p_m) \leq p_{m+1}$. We prove (b) by induction. $n = 1$ is the definition of Z . Now suppose that for $n \geq 2$

$$Z_{n-1}(x) = p V^{(n-1)} x \otimes \mathbb{1} V^{(n-1)*} |_{\mathfrak{h}} = Z^{n-1}(x).$$

Then we can use (2) of Lemma 2.1 in the form $pV = pV(p \otimes \mathbb{1})$ to get

$$\begin{aligned} Z_n(x) &= p V^{(n)} x \otimes \mathbb{1} V^{(n)*} |_{\mathfrak{h}} \\ &= p V[(V^{(n-1)} \otimes \mathbb{1})((x \otimes \mathbb{1}) \otimes \mathbb{1})(V^{(n-1)*} \otimes \mathbb{1})] V^* |_{\mathfrak{h}} \\ &= p V(p \otimes \mathbb{1})[\dots](p \otimes \mathbb{1}) V^* |_{\mathfrak{h}} \\ &= p V(Z^{n-1}(x) \otimes \mathbb{1}) V^* |_{\mathfrak{h}} = Z^n(x) \end{aligned}$$

To prove (c) we note that (2) of Lemma 2.1 in the form $V(\mathfrak{h}^\perp \otimes \mathcal{P}) \perp \mathfrak{h}$ can be iterated to yield

$$V^{(\ell)}(\mathfrak{h}^\perp \otimes \mathcal{P}^{\otimes \ell}) \perp \mathfrak{h}.$$

for all $\ell \in \mathbb{N}$. Hence for all $\ell \leq m$ we find, by applying $V^{(m-\ell)}$,

$$V^{(m)}(\mathfrak{h}^\perp \otimes \mathcal{P}^{\otimes m}) \perp V^{(m-\ell)}(\mathfrak{h} \otimes \mathcal{P}^{\otimes (m-\ell)})$$

which implies $p_m V^{(m)}(p^\perp \otimes \mathbb{1}) = 0$.

Together with $q_m = \theta^m(p) = V^{(m)} p \otimes \mathbb{1} V^{(m)*}$ we obtain

$$p_m V^{(m)} = p_m V^{(m)} p \otimes \mathbb{1} = p_m q_m V^{(m)} = q_m V^{(m)}.$$

The iterative definition of the $V^{(m)}$ shows that their ranges do not increase if m increases. Hence also $p_m V^{(n)} = q_m V^{(n)}$ whenever $m \leq n$.

To get (d) we start from the definition of Z_{n-m} ,

$$p J^{(n-m)}(x) p = Z_{n-m}(x) p, \quad (m \leq n),$$

and apply θ^m to get

$$q_m J^{(n)}(x) q_m = \theta^m(Z_{n-m}(x) p) = J^{(m)}(Z_{n-m}(x))$$

Now because of (b) and (c) we can replace Z_{n-m} by Z^{n-m} and q_m by p_m . □

The following observation, also noted in [10], is crucial for our approach. Bhat remarks in [10], p.562, in this context: ‘Roughly speaking there is also an additive structure when we deal with general quantum dynamical semigroups ... A detailed study of such systems is yet to be undertaken.’ The work presented in this paper goes into this direction.

Proposition 2.5. Let $(\mathcal{H}, V, \mathfrak{h})$ be a process. Then

$$\mathcal{E} := \mathfrak{h}_1 \ominus \mathfrak{h}_0 = \overline{\text{span}}(\mathfrak{h}, V(\mathfrak{h} \otimes \mathcal{P})) \ominus \mathfrak{h}$$

is a wandering subspace, i.e., $V_\alpha \mathcal{E} \perp V_\beta \mathcal{E}$ if $\alpha \neq \beta$, and

$$\mathcal{H} = \mathfrak{h} \oplus \bigoplus_{\alpha \in F_d^+} V_\alpha \mathcal{E}, \quad \mathfrak{h}_n = \mathfrak{h} \oplus \bigoplus_{|\alpha| < n} V_\alpha \mathcal{E}$$

Proof. To prove that $\mathcal{E}$ is wandering it is enough to show that $\mathcal{E} \perp V_\alpha \mathcal{E}$ for all $\alpha \neq 0$. Suppose $\alpha \neq 0$. Because $\mathcal{E} \perp \mathfrak{h}$ we conclude by (2) of Lemma 2.1 that $V_\beta \mathcal{E} \perp \mathfrak{h}$ for all $\beta \in F_d^+$. In particular $V_\alpha \mathcal{E} \perp \mathfrak{h}$. Writing $\alpha = k\beta$ with $k \in \{1, \dots, d\}$ and $\beta \in F_d^+$ we find $V_\alpha \mathcal{E} \perp V_k \mathfrak{h}$. Finally if $k' \in \{1, \dots, d\}$ but $k' \neq k$ then $V_\alpha \mathcal{E} \perp V_{k'} \mathcal{H}$ because V_k and $V_{k'}$ have orthogonal ranges. Putting it all together we have $\mathcal{E} \perp V_\alpha \mathcal{E}$. The other assertions are now immediate. $\square$

The following results indicate how the V_α are related to the quantum physical behaviour of the process. For this we apply the standard interpretations of quantum physics to the mathematical objects. Suppose $X : \mathcal{H} \rightarrow \mathcal{H}$ and $Y : \mathcal{P} \rightarrow \mathcal{P}$ are linear operators. Then for $n \in \mathbb{N}_0$ and $m \in \mathbb{N}$ we have the following linear operators on $\mathcal{H}$

$$\begin{aligned} X_n &:= V^{(n)} X \otimes \bigotimes_{i=1}^n \mathbb{1}_{\mathcal{P}} V^{(n)*}, \\ Y_m &:= V^{(m)} \mathbb{1}_{\mathcal{H}} \otimes \bigotimes_{i=1}^{m-1} \mathbb{1}_{\mathcal{P}} \otimes Y V^{(m)*} \end{aligned}$$

(empty tensor products to be omitted for $n = 0$ and $m = 1$). We refer to these operators as observables.

Proposition 2.6. Suppose that the process $(\mathcal{H}, V, \mathfrak{h})$ is unital. Then the observables Y_m commute with each other and Y_m commutes with X_n whenever $m \leq n$.

Suppose further that $Y\epsilon_j = j\epsilon_j$ for the orthonormal basis $(\epsilon_j)_{j=1}^d$ of $\mathcal{P}$ which we use to define the isometries V_j . If the process is prepared in a vector state given by a unit vector η in the range of V_α so that the m -th letter in α is equal to k then the measurement of Y_m yields the outcome k with certainty.

Proof. For $1 \leq m \leq n$ we define

$$Y_{m,n} := V^{(n)} \mathbb{1}_{\mathcal{H}} \otimes \bigotimes_{i=1}^{m-1} \mathbb{1}_{\mathcal{P}} \otimes Y \otimes \bigotimes_{i=m+1}^n \mathbb{1}_{\mathcal{P}} V^{(n)*}$$

(empty tensor products omitted in the cases $m = 1$ and $m = n$). Because the process is unital, V is a row unitary and in particular $V \mathbb{1}_{\mathcal{H}} \otimes \mathbb{1}_{\mathcal{P}} V^* = \mathbb{1}_{\mathcal{H}}$. It follows that $Y_m = Y_{m,n}$ whenever $m \leq n$. Writing Y_m in this way it becomes obvious that it commutes with X_n and Y_n .

Let P_{ϵ_k} be the orthogonal projection onto the one-dimensional subspace $\mathbb{C}\epsilon_k$ of $\mathcal{P}$. Then

$$\langle \eta, V^{(m)} \mathbb{1}_{\mathcal{H}} \otimes \bigotimes_{i=1}^{m-1} \mathbb{1}_{\mathcal{P}} \otimes P_{\epsilon_k} V^{(m)*} \eta \rangle = \langle V^{(m)*} \eta, \mathbb{1}_{\mathcal{H}} \otimes \bigotimes_{i=1}^{m-1} \mathbb{1}_{\mathcal{P}} \otimes P_{\epsilon_k} V^{(m)*} \eta \rangle$$

is the probability that a measurement of Y_m yields the result k , according to the standard rules of quantum mechanics. Now suppose that the process is prepared in a vector state with a unit vector η in the range of V_α , i.e., $\eta = V_\alpha \xi$ for some $\xi \in \mathcal{H}$, so that the m -th letter in α is equal to k . Then from

$$V^{(m)*} \eta = \sum_{|\beta|=m} V_\beta^* V_\alpha \xi \otimes \epsilon_\beta$$

(with ϵ_β for $\beta = \beta_1 \dots \beta_m$ is short for $\epsilon_{\beta_1} \otimes \dots \otimes \epsilon_{\beta_m} \in \bigotimes_1^m \mathcal{P}$) we see that only the terms in the sum with $\beta_m = k$ can be non-zero and hence the application of P_{ϵ_k} at the m -th copy of $\mathcal{P}$ always acts identical on all the non-zero terms. So the probability above is the squared length of a unit vector, i.e., it is equal to 1. $\square$

This result gives an operational meaning to the words $\alpha \in F_d^+$ by identifying them with measurement protocols for certain observables. We can think of measuring $Y_1, \dots, Y_n$ as performing a certain type Y of measurement at the consecutive times $1, \dots, n$ and the commutation properties proved above ensure that these measurements can be performed without perturbing the system (non-demolition measurements). For example with $x \in \mathcal{B}(\mathfrak{h})$ it makes sense to consider $J^{(n)}(x) = X_n$ for $X = xp_{\mathfrak{h}}$ conditioned on $Y_1, \dots, Y_n$ (conditioning in the sense of classical probability theory). Such schemes and their application to quantum filtering and quantum control have been pioneered by Belavkin [6, 7] and this is a rapidly developing interdisciplinary field of study. We mention the recent introductory surveys [12, 13] which give many references, the latter focusing on discretized models and containing constructions similar to the one above ([13], Section 2.5). It is of course interesting to prepare the system in other states where the outcomes are not deterministic but only statistical information can be obtained, see for example [28] which uses discretization of continuous processes and thus can be directly connected with the approach here. For a unital process the Hilbert space $\mathcal{H}$ is always the orthogonal sum of the ranges of the V_α for all α with a given length, so in principle the problem can be dealt with for an arbitrary vector state by decomposing the state vector with respect to such an orthogonal sum and then using Proposition 2.6. This may not always be the most practical path to follow for data given in a different way but it should be of theoretical significance. In this place we only want to indicate the relevance of the V_α for applications and so we don't go into further details here.

3. REPRESENTATIONS OF STRUCTURE MAPS

If for the multi-variable systems introduced in Section 1 the spaces are Hilbert spaces and the linear maps are contractions between these Hilbert spaces then we want to think of them as appearing inside the weak Markov processes introduced in Section 2. The following definitions give a precise meaning to that. Then we justify the definitions by discussing how the represented structure maps can help us to understand the properties of the process.

Definition 3.1. A *representation of an input pair* (A, B) (with column contractions $A : \mathcal{X} \rightarrow \bigoplus_1^d \mathcal{X}$ and $B : \mathcal{U} \rightarrow \bigoplus_1^d \mathcal{X}$) is given by a process $(\mathcal{H}, V, \mathfrak{h})$ such that $\mathcal{X} = \mathfrak{h}$ and $A = V^*|_{\mathfrak{h}}$ together with an isometry $i_0 : \mathcal{U} \rightarrow \mathcal{E}$ such that, with $\mathcal{U}_0 := i_0(\mathcal{U})$ the represented input space, we have $B = V^*|_{\mathcal{U}_0} i_0$.

A *representation of an output pair* (A, C) (with column contractions $A : \mathcal{X} \rightarrow \bigoplus_1^d \mathcal{X}$ and $C : \mathcal{X} \rightarrow \mathcal{Y}$) is given by a process $(\mathcal{H}, V, \mathfrak{h})$ such that $\mathcal{X} = \mathfrak{h}$ and $A = V^*|_{\mathfrak{h}}$ together with an isometry $j_0 : \mathcal{Y} \rightarrow \mathfrak{h}_1 = \mathfrak{h} \oplus \mathcal{E}$ such that the represented output space $\mathcal{Y}_0 := j_0(\mathcal{Y})$ is wandering for V and we have $C = j_0^* P_{\mathcal{Y}_0}|_{\mathfrak{h}}$.

A *representation of the structure maps* (A, B, C, D) , where $D : \mathcal{U} \rightarrow \mathcal{Y}$, is given by representations of (A, B) as an input pair and (A, C) as an output pair as above with the same process $(\mathcal{H}, V, \mathfrak{h})$ such that $D = j_0^* P_{\mathcal{Y}_0}|_{\mathcal{U}_0} i_0$.

In the following we shall often suppress the isometries i_0 and j_0 and treat them as identifications whenever this simplifies the notation. Note that a represented input space $\mathcal{U}_0$ is always wandering because by definition it is a subspace of the wandering subspace $\mathcal{E}$, see Proposition 2.5. An interesting example for a represented output space is

$$\mathcal{E}_* := \mathfrak{h}_1 \ominus V(\mathfrak{h} \otimes \mathcal{P}) = \mathcal{H} \ominus V(\mathcal{H} \otimes \mathcal{P}) = \ker V^*$$

which is the wandering subspace arising from the Wold decomposition of the row isometry V , see [33].

Let us write $\mathcal{U}_\beta$ for $V_\beta \mathcal{U}_0$ and $\mathcal{Y}_\alpha$ for $V_\alpha \mathcal{Y}_0$, further we denote by $\mathcal{U}_+$ the closed linear span of all $\mathcal{U}_\alpha$, by $\mathcal{Y}_+$ the closed linear span of all $\mathcal{Y}_\alpha$ (with $\alpha \in F_d^+$). With suitable identifications we can think of $\mathcal{U}_+$ as an orthogonal direct sum of copies of the input space $\mathcal{U}$, similarly $\mathcal{Y}_+$ is an orthogonal direct sum of copies of the output space $\mathcal{Y}$. We sometimes refer to these copies as α -translated input resp. output spaces.

Representations of structure maps as above are always *causal* in the sense that $\mathcal{Y}_0 \perp V_\beta \mathcal{U}_0 = \mathcal{U}_\beta$ for all $\beta \in F_d^+$ with $|\beta| \geq 1$. More general it follows further that $\mathcal{Y}_\alpha \perp \mathcal{U}_\beta$ whenever $|\alpha| < |\beta|$ which we also refer to as causality.

Note that from a representation with $A = (A_1, \dots, A_d)^t$ we get a Kraus decomposition for the transition operator of the process, namely

$$Z(x) = A^* x A = \sum_{k=1}^d A_k^* x A_k,$$

(limit in the strong operator topology if $d = \infty$).

Given a process $(\mathcal{H}, V, \mathfrak{h})$ any subspace of $\mathcal{E}$ interpreted as a represented input space gives rise to a representation of an input pair (A, B) and any wandering subspace of $\mathfrak{h}_1$ interpreted as a represented output space gives rise to a representation of an output pair (A, C) and together we have a representation of structure maps (A, B, C, D) .

Proposition 3.2. Given a representation of structure maps (A, B, C, D) by a process $(\mathcal{H}, V, \mathfrak{h})$ let $\tilde{\xi} = \xi \oplus \bigoplus V_\alpha \eta_\alpha$ be an element of $\mathcal{H}$, with $\xi \in \mathfrak{h}$ and $\eta_\alpha \in \mathcal{U}_0$. If (for all words α)

$$\begin{aligned} x(0) &:= \xi, & x(\alpha) &:= P_{\mathfrak{h}} V_\alpha^* \tilde{\xi} \\ u(\alpha) &:= \eta_\alpha & (\Rightarrow V_\alpha u(\alpha) &= P_{\mathcal{U}_\alpha} \tilde{\xi}) \end{aligned}$$

then we have (for all words α and generators $k = 1, \dots, d$)

$$x(\alpha k) = A_k x(\alpha) + B_k u(\alpha).$$

If further $y(\alpha) := P_{\mathcal{Y}_0} V_\alpha^* \tilde{\xi}$ ($\Rightarrow V_\alpha y(\alpha) = P_{\mathcal{Y}_\alpha} \tilde{\xi}$) then we have

$$y(\alpha) = C x(\alpha) + D u(\alpha).$$

Hence we have a noncommutative Fornasini-Marchesini system (compare Section 1).

Proof. The first assertion follows from

$$\begin{aligned} x(\alpha k) &= P_{\mathfrak{h}} V_{\alpha k}^* \tilde{\xi} = P_{\mathfrak{h}} V_k^* V_\alpha^* \tilde{\xi} = P_{\mathfrak{h}} V_k^* P_{\mathfrak{h} \oplus \mathcal{U}_0} V_\alpha^* \tilde{\xi} \\ &= V_k^* P_{\mathfrak{h}} V_\alpha^* \tilde{\xi} + V_k^* \eta_\alpha = A_k x(\alpha) + B_k u(\alpha). \end{aligned}$$

Using causality we find that

$$\begin{aligned} y(\alpha) &= P_{\mathcal{Y}_0} V_\alpha^* \tilde{\xi} = P_{\mathcal{Y}_0} P_{\mathfrak{h} \oplus \mathcal{U}_0} V_\alpha^* \tilde{\xi} \\ &= P_{\mathcal{Y}_0} [x(\alpha) + u(\alpha)] = C x(\alpha) + D u(\alpha). \end{aligned}$$

□

Note that the proof uses causality but does not use the fact that $\mathcal{Y}_0$ is wandering. Hence the result is valid for generalized (non-wandering but causal) output spaces.

With the maximal choice $\mathcal{U}_0 = \mathcal{E}$ we have for any vector $\tilde{\xi} \in \mathcal{H}$ that

$$P_{V_\alpha \mathcal{H}} \tilde{\xi} = P_{V_\alpha \mathfrak{h}} \tilde{\xi} \oplus \bigoplus_{\beta \in F_d^+} P_{V_\alpha \mathcal{U}_\beta} \tilde{\xi} = V_\alpha x(\alpha) \oplus \bigoplus_{\beta \in F_d^+} V_{\alpha\beta} u(\alpha\beta),$$

which is relevant information with respect to Proposition 2.6 above.

Let us now work out the observability map $\mathcal{O}_{C,A}$ and the transfer function $\mathcal{T}$ (already introduced in Section 1) for represented structure maps. If we identify $\mathcal{Y}_\alpha \simeq_\alpha \mathcal{Y}$ then

$$P_{\mathcal{Y}_\alpha}|_{\mathfrak{h}} = V_\alpha P_{\mathcal{Y}_0} V_\alpha^*|_{\mathfrak{h}} \simeq_\alpha C A^\alpha$$

in other words, we obtain the α -entry $C A^\alpha$ of the observability map $\mathcal{O}_{C,A}$ by projecting the representation $\mathfrak{h}$ of the internal space $\mathcal{X}$ to the α -translated output space $\mathcal{Y}_\alpha$.

Similarly with $\mathcal{Y}_\alpha \simeq_\alpha \mathcal{Y}$ and $\mathcal{U}_0 \simeq \mathcal{U}$ we obtain

$$P_{\mathcal{Y}_\alpha}|_{\mathcal{U}_0} = V_\alpha P_{\mathcal{Y}_0} V_\alpha^*|_{\mathcal{U}_0} \simeq_\alpha \begin{cases} D & \text{if } \alpha = 0 \\ C B_\alpha & \text{if } |\alpha| = 1 \\ C A_{\alpha_r} \dots A_{\alpha_2} B_{\alpha_1} & \text{if } \alpha = \alpha_1 \dots \alpha_r, r = |\alpha| \geq 2 \end{cases}$$

In other words, we obtain the α -coefficient $\mathcal{T}^\alpha$ in the formal power series expansion of the transfer function $\mathcal{T}$ (already introduced in Section 1) by projecting the representation $\mathcal{U}_0$ of the input space $\mathcal{U}$

to the α -translated output space $\mathcal{Y}_\alpha$. Causality implies that further, with identifications $\mathcal{Y}_\alpha \simeq_\alpha \mathcal{Y}$ and $\mathcal{U}_\beta \simeq_\beta \mathcal{U}$,

$$P_{\mathcal{Y}_\alpha|_{\mathcal{U}_\beta}} \simeq_{\alpha,\beta} \begin{cases} \mathcal{T}^\sigma & \text{if } \alpha = \beta\sigma \\ 0 & \text{otherwise} \end{cases}$$

This pattern in the operator-valued kernel $(P_{\mathcal{Y}_\alpha|_{\mathcal{U}_\beta}})_{\alpha,\beta \in F_d^+}$ describes what is called a multi-analytic kernel. It is an operator-valued matrix describing the contraction $P_{\mathcal{Y}_+|_{\mathcal{U}_+}}$ for the orthogonal decompositions with respect to the translated output resp. input spaces. It follows from this description that this contraction intertwines the row shifts on $\bigoplus_\beta \mathcal{U}_\beta$ and $\bigoplus_\alpha \mathcal{Y}_\alpha$ which are obtained by restricting the row isometry V . This is the defining property of a multi-analytic operator and we have verified it for the contraction $P_{\mathcal{Y}_+|_{\mathcal{U}_+}}$. Multi-analytic operators have been introduced and studied by Popescu [34, 36], the situation above involving transfer functions for pairs of wandering subspaces is worked out in more detail in [25].

We can think of it as a kind of multiplication by the transfer function $\mathcal{T}$. In fact for $d = 1$ it is literally multiplication by an ordinary analytic function. But for all $d \geq 1$, for a noncommutative Fornasini-Marchesini system, we can still verify the following analogy: If we introduce formal power series for the entries $y(\alpha)$ and $u(\beta)$ (see Proposition 3.2)

$$\begin{aligned} \hat{u}(z) &= \hat{u}(z_1, \dots, z_d) = \sum_\beta u(\beta) z^\beta \\ \hat{y}(z) &= \hat{y}(z_1, \dots, z_d) = \sum_\alpha y(\alpha) z^\alpha \end{aligned}$$

then, with the initial condition $x(0) = 0$, we have the input-output relation

$$\hat{y}(z) = \mathcal{T}(z) \hat{u}(z).$$

Recall the convention $z^\alpha = z_{\alpha_r} \dots z_{\alpha_1}$ for $\alpha = \alpha_1 \dots \alpha_r$ which gives the multiplication rule $z^\sigma z^\beta = z^{\beta\sigma}$.

4. CATEGORIES OF PROCESSES AND γ -CASCADES

We introduce additional concepts with the intention to describe substructures of processes. We say that $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g})$ is a *subprocess* of the process $(\mathcal{H}, V, \mathfrak{h})$ if $\mathfrak{g}$ is a closed subspace of $\mathfrak{h}$ which is co-invariant for V and $V^\mathcal{G} = V|_{\mathcal{G}}$ where $\mathcal{G} = \overline{\text{span}}\{V_\alpha \mathfrak{g} : \alpha \in F_d^+\}$. Note that $\mathfrak{g}$ is also co-invariant for $V^\mathcal{G}$ and $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g})$ is a process in its own right.

Given a subprocess $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g})$ of a process $(\mathcal{H}, V, \mathfrak{h})$ we can form the *quotient process*

$$(\mathcal{H}, V, \mathfrak{h})/(\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) := (\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$$

where $\mathfrak{k} := \mathfrak{h} \ominus \mathfrak{g}$, $\mathcal{K} := \overline{\text{span}}\{V_\alpha \mathfrak{k} : \alpha \in F_d^+\}$, $V^\mathcal{K} := V|_{\mathcal{K}}$. Let us check that $(\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$ is a process. Indeed, because $\mathfrak{g}$ is co-invariant for V we see that $\mathcal{K}$ is contained in $\mathcal{H} \ominus \mathfrak{g}$. Hence $\mathcal{K} \ominus \mathfrak{k}$ is contained in $\mathcal{H} \ominus \mathfrak{h}$ which is a V -invariant subspace orthogonal to $\mathfrak{k}$. Hence $\mathfrak{k}$ is co-invariant for $V^\mathcal{K}$ which proves our claim. Note that in general $\mathfrak{k}$ is not co-invariant for V . Considering adjoints we see that $V^{\mathcal{G}*} = V^*|_{\mathcal{G}}$ but only $V^{\mathcal{K}*} = (P_{\mathcal{K}} \otimes \mathbb{1}_{\mathcal{P}}) V^*|_{\mathcal{K}}$. So we need to distinguish carefully between subprocesses and quotient processes.

It is convenient to reformulate these concepts within a *category of processes* which we define now. The objects of the category are the processes with a common multiplicity space $\mathcal{P}$. A *morphism* from $(\mathcal{R}, V^\mathcal{R}, \mathfrak{r})$ to $(\mathcal{S}, V^\mathcal{S}, \mathfrak{s})$ is a contraction $t : \mathfrak{r} \rightarrow \mathfrak{s}$ which intertwines the adjoints of the row isometries, i.e.

$$V^{\mathcal{S}*} t = (t \otimes \mathbb{1}_{\mathcal{P}}) V^{\mathcal{R}*}|_{\mathfrak{r}}.$$

(or written differently: $A_k^{\mathcal{S}} t = t A_k^{\mathcal{R}}$ for $k = 1, \dots, d$ and $A_k^{\mathcal{S}} = V_k^{\mathcal{S}*}|_{\mathfrak{s}}$, $A_k^{\mathcal{R}} = V_k^{\mathcal{R}*}|_{\mathfrak{r}}$). Composition of morphisms is given by composition of operators, the identity morphism is given by the identity operator.

We add some immediate observations justifying this definition. First note that a morphism t is an isomorphism if and only if t is unitary and we get a reasonable meaning for two processes to be isomorphic. In the following we often identify isomorphic processes.

Also note that if we have a morphism given by a contraction $t : \mathfrak{r} \rightarrow \mathfrak{s}$ then t can always be extended to a contraction $T : \mathcal{R} \rightarrow \mathcal{S}$ such that $T|_{\mathfrak{r}} = t$ and $\|T\| = \|t\|$ and

$$V^{\mathcal{S}*}T = (T \otimes \mathbb{1}_{\mathcal{P}})V^{\mathcal{R}*},$$

this is nothing but the commutant lifting theorem in the version of Popescu [35]. We call T an *extended morphism* associated to t . We can also think of a morphism as the class of all extended morphisms associated to t , this attaches a global interpretation to it.

Finally it follows from $V^{\mathcal{S}*}t = (t \otimes \mathbb{1}_{\mathcal{P}})V^{\mathcal{R}*}|_{\mathfrak{r}}$ that $t(\mathfrak{r})$ is co-invariant for $V^{\mathcal{S}}$, and hence from a morphism t from $(\mathcal{R}, V^{\mathcal{R}}, \mathfrak{r})$ to $(\mathcal{S}, V^{\mathcal{S}}, \mathfrak{s})$, by defining $\mathfrak{g}$ to be the closure of $t(\mathfrak{r})$ and $V^{\mathcal{G}} := V^{\mathcal{S}}|_{\mathcal{G}}$, we always obtain a subprocess $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ of $(\mathcal{S}, V^{\mathcal{S}}, \mathfrak{s})$.

In fact, we can reformulate subprocesses and quotient processes in terms of morphisms as follows. If $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ is a subprocess of a process $(\mathcal{H}, V, \mathfrak{h})$ then $\mathbb{1}_{\mathfrak{g}}$, interpreted as an embedding of $\mathfrak{g}$ into $\mathfrak{h}$, is an isometric morphism:

$$V^*\mathbb{1}_{\mathfrak{g}} = (\mathbb{1}_{\mathfrak{g}} \otimes \mathbb{1}_{\mathcal{P}})V^{\mathcal{G}*}|_{\mathfrak{g}}$$

Conversely, given an isometric morphism t from a process $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ to a process $(\mathcal{H}, V, \mathfrak{h})$ we can interpret t as an embedding $\mathbb{1}_{\mathfrak{g}}$ and the properties of a morphism ensure that $\mathfrak{g}$ is co-invariant for V . With an extended morphism $\mathbb{1}_{\mathcal{G}}$, interpreted as an embedding of $\mathcal{G}$ into $\mathcal{H}$, we can also arrange that $V^{\mathcal{G}} = V|_{\mathcal{G}}$ (by uniqueness of a minimal isometric dilation up to unitary equivalence). So we get a subprocess.

For the corresponding quotient process $(\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$ we can check that the orthogonal projection $P_{\mathfrak{k}} : \mathfrak{h} \rightarrow \mathfrak{k}$ is a coisometric morphism:

$$V^{\mathcal{K}*}P_{\mathfrak{k}} = (P_{\mathfrak{k}} \otimes \mathbb{1}_{\mathcal{P}})V^*|_{\mathfrak{h}}$$

Conversely, if we have a coisometric morphism t from $(\mathcal{H}, V, \mathfrak{h})$ to a process $(\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$ then we can interpret t as an orthogonal projection $P_{\mathfrak{k}} : \mathfrak{h} \rightarrow \mathfrak{k}$, then use the properties of a morphism to check that $\mathfrak{g} := \mathfrak{h} \ominus \mathfrak{k}$ is co-invariant for V , giving rise to a subprocess so that the corresponding quotient process is the process $(\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$ we started from. Note that $P_{\mathfrak{g}}$ and $\mathbb{1}_{\mathfrak{k}}$ are not morphisms, in general.

The image of $\mathbb{1}_{\mathfrak{g}}$ is equal to the kernel of $P_{\mathfrak{k}}$ and we can proceed to give a concise description of the situation as a short exact sequence of processes:

$$0 \longrightarrow (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \xrightarrow{\mathbb{1}_{\mathfrak{g}}} (\mathcal{H}, V, \mathfrak{h}) \xrightarrow{P_{\mathfrak{k}}} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) \longrightarrow 0$$

Here 0 stands for the process on the 0-dimensional space which takes the role of a zero object in our category. In the following we suppress the embeddings in the notation whenever this is convenient.

Lemma 4.1. Given a short exact sequence of processes

$$0 \longrightarrow (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \xrightarrow{\mathbb{1}_{\mathfrak{g}}} (\mathcal{H}, V, \mathfrak{h}) \xrightarrow{P_{\mathfrak{k}}} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) \longrightarrow 0$$

then for the relative position of the wandering subspaces

$$\begin{aligned} \mathcal{E}^{\mathfrak{g}} &:= \overline{\text{span}}(\mathfrak{g}, V^{\mathcal{G}}(\mathfrak{g} \otimes \mathcal{P})) \ominus \mathfrak{g} \\ \mathcal{E}_*^{\mathfrak{k}} &:= \overline{\text{span}}(\mathfrak{k}, V^{\mathcal{K}}(\mathfrak{k} \otimes \mathcal{P})) \ominus V^{\mathcal{K}}(\mathfrak{k} \otimes \mathcal{P}) \end{aligned}$$

we obtain

$$P_{\mathcal{G}}|_{\mathcal{E}_*^{\mathfrak{k}}} = P_{\mathcal{E}^{\mathfrak{g}}}|_{\mathcal{E}_*^{\mathfrak{k}}}, \quad P_{\mathcal{K}}|_{\mathcal{E}^{\mathfrak{g}}} = P_{\mathcal{E}_*^{\mathfrak{k}}}|_{\mathcal{E}^{\mathfrak{g}}} = (P_{\mathcal{E}^{\mathfrak{g}}}|_{\mathcal{E}_*^{\mathfrak{k}}})^*$$

More general, for all $\alpha \in F_d^+$

$$\begin{aligned} P_{\mathcal{G}}|_{V_{\alpha}\mathcal{E}_*^{\mathfrak{k}}} &= P_{V_{\alpha}\mathcal{E}^{\mathfrak{g}}}|_{V_{\alpha}\mathcal{E}_*^{\mathfrak{k}}} = V_{\alpha}P_{\mathcal{E}^{\mathfrak{g}}}|_{\mathcal{E}_*^{\mathfrak{k}}}V_{\alpha}^*|_{V_{\alpha}\mathcal{E}_*^{\mathfrak{k}}} \\ P_{\mathcal{K}}|_{V_{\alpha}\mathcal{E}^{\mathfrak{g}}} &= P_{V_{\alpha}\mathcal{E}_*^{\mathfrak{k}}}|_{V_{\alpha}\mathcal{E}^{\mathfrak{g}}} = V_{\alpha}P_{\mathcal{E}_*^{\mathfrak{k}}}|_{\mathcal{E}^{\mathfrak{g}}}V_{\alpha}^*|_{V_{\alpha}\mathcal{E}^{\mathfrak{g}}} \end{aligned}$$

Proof. We have $\mathcal{E}_*^{\mathfrak{k}} \subset \mathfrak{h} \oplus \mathcal{E}$ where $\mathcal{E} := \overline{\text{span}}(\mathfrak{h}, V(\mathfrak{h} \otimes \mathcal{P})) \ominus \mathfrak{h}$. Because $V^*\mathfrak{h} \oplus \mathcal{E} = \mathfrak{h} \otimes \mathcal{P}$ we conclude that $V^*\mathcal{E}_*^{\mathfrak{k}} \subset \mathfrak{h} \otimes \mathcal{P}$. But by the definition of $\mathcal{E}_*^{\mathfrak{k}}$ we also have

$$(P_{\mathcal{K}} \otimes \mathbb{1}_{\mathcal{P}})V^*\mathcal{E}_*^{\mathfrak{k}} = V^{\mathcal{K}*}\mathcal{E}_*^{\mathfrak{k}} = \{0\}$$

and it follows that $V^* \mathcal{E}_*^\mathfrak{k} \subset \mathfrak{g} \otimes \mathcal{P}$. Hence if $\xi \in \mathcal{E}^\mathfrak{g}$, $\eta \in \mathcal{E}_*^\mathfrak{k}$ and $\alpha \neq 0$ then $V_\alpha^* \eta \subset \mathfrak{g}$ and

$$\langle V_\alpha \xi, \eta \rangle = \langle \xi, V_\alpha^* \eta \rangle = 0$$

and it follows that $\mathcal{E}_*^\mathfrak{k} \perp V_\alpha \mathcal{E}^\mathfrak{g}$ for all $\alpha \neq 0$. Obviously also $\mathcal{E}_*^\mathfrak{k} \perp \mathfrak{g}$ and so $P_{\mathcal{E}^\mathfrak{g}}|_{\mathcal{E}_*^\mathfrak{k}} = P_{\mathcal{G}}|_{\mathcal{E}_*^\mathfrak{k}}$ which is the first formula we intended to prove.

It is clear that $(P_{\mathcal{E}^\mathfrak{g}}|_{\mathcal{E}_*^\mathfrak{k}})^* = P_{\mathcal{E}_*^\mathfrak{k}}|_{\mathcal{E}^\mathfrak{g}}$. To get the equality $P_{\mathcal{K}}|_{\mathcal{E}^\mathfrak{g}} = P_{\mathcal{E}_*^\mathfrak{k}}|_{\mathcal{E}^\mathfrak{g}}$ we argue as follows. Consider $\mathcal{E}^\mathfrak{k} := \overline{\text{span}}(\mathfrak{k}, V(\mathfrak{k} \otimes \mathcal{P})) \ominus \mathfrak{k}$. Because $\mathcal{E}^\mathfrak{k} \perp \mathfrak{h}$ it follows that $\mathcal{E}^\mathfrak{k} \subset \mathcal{E}$ and because $\mathcal{E}$ is wandering for V we conclude that $V_\alpha \mathcal{E}^\mathfrak{k} \perp \mathfrak{h} \oplus \mathcal{E}$ if $\alpha \neq 0$ and hence

$$P_{\mathcal{K}}(\mathfrak{h} \oplus \mathcal{E}) = \mathfrak{k} \oplus \mathcal{E}^\mathfrak{k}.$$

Now, because $\mathcal{E}^\mathfrak{g} \subset \mathfrak{h} \oplus \mathcal{E}$, we can compute

$$P_{\mathcal{K}}|_{\mathcal{E}^\mathfrak{g}} = P_{\mathcal{K}}P_{\mathfrak{h} \oplus \mathcal{E}}|_{\mathcal{E}^\mathfrak{g}} = P_{\mathfrak{k} \oplus \mathcal{E}^\mathfrak{k}}|_{\mathcal{E}^\mathfrak{g}} = P_{\mathcal{E}_*^\mathfrak{k}}|_{\mathcal{E}^\mathfrak{g}}.$$

For the last equality above note that $\mathcal{E}_*^\mathfrak{k} = (\mathfrak{k} \oplus \mathcal{E}^\mathfrak{k}) \ominus V(\mathfrak{k} \otimes \mathcal{P})$ but $\mathcal{E}^\mathfrak{g} \subset \overline{\text{span}}\{g, V(\mathfrak{g} \otimes \mathcal{P})\} \perp V(\mathfrak{k} \otimes \mathcal{P})$.

Because $\mathcal{E}^\mathfrak{g}$ and $\mathcal{E}_*^\mathfrak{k}$ are wandering it is now immediate that

$$V_\alpha \mathcal{E}^\mathfrak{g} \perp V_\beta \mathcal{E}_*^\mathfrak{k} \quad \text{if } \alpha \neq \beta.$$

To get the first line of the general formulas we only have to note additionally that always $V_\beta \mathcal{E}_*^\mathfrak{k} \perp \mathfrak{g}$. To get the second line we have to prove that additionally

$$V_\alpha \mathcal{E}^\mathfrak{g} \perp \mathcal{K} \ominus \bigoplus_{\beta \in F_d^+} V_\beta \mathcal{E}_*^\mathfrak{k} =: \mathcal{K}_{res} \quad (\text{residual part}).$$

Indeed, if $\alpha = 0$ this follows from $P_{\mathcal{K}}|_{\mathcal{E}^\mathfrak{g}} = P_{\mathcal{E}_*^\mathfrak{k}}|_{\mathcal{E}^\mathfrak{g}}$ shown above. Then for other α reduce it to the previous case by using the fact that $V^\mathcal{K}|_{\mathcal{K}_{res}}$ is a row unitary and hence $V^* \mathcal{K}_{res} = V^\mathcal{K} \mathcal{K}_{res} \subset \mathcal{K}_{res} \otimes \mathcal{P}$. $\square$

These observations suggest the following construction which allows us to classify the processes which can be obtained by short exact sequences, given the subprocess and the quotient process.

Definition 4.2. Given processes $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g})$ and $(\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$ (with a common multiplicity space $\mathcal{P}$) and any contraction $\gamma : \mathcal{E}_*^\mathfrak{k} \rightarrow \mathcal{E}^\mathfrak{g}$ we define the γ -cascade

$$(\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) \triangleleft_\gamma (\mathcal{K}, V^\mathcal{K}, \mathfrak{k}) := (\mathcal{H}, V, \mathfrak{h})$$

where

$$\begin{aligned} \mathfrak{h} &:= \mathfrak{g} \oplus \mathfrak{k}, \\ \mathcal{H} &:= \mathfrak{g} \oplus \mathcal{K} \oplus \bigoplus_{\alpha \in F_d^+} (\mathcal{D}_{\gamma^*})_\alpha, \\ V &:= \begin{cases} (\mathbb{1}_{\mathfrak{g}} \oplus \begin{pmatrix} \gamma^* \\ D_{\gamma^*} \end{pmatrix}) V^\mathcal{G} & \text{on } \mathfrak{g} \\ V^\mathcal{K} & \text{on } \mathcal{K} \\ \text{canonical row shift} & \text{on } \bigoplus_{\alpha \in F_d^+} (\mathcal{D}_{\gamma^*})_\alpha \end{cases} \end{aligned}$$

We add the following explanations for this definition. $D_{\gamma^*} := \sqrt{\mathbb{1} - \gamma\gamma^*}$ is the defect operator for γ^* and $\mathcal{D}_{\gamma^*}$, the closure of its range, is the defect space. Then $\bigoplus_{\alpha \in F_d^+} (\mathcal{D}_{\gamma^*})_\alpha$ is the orthogonal sum of a family of copies of $\mathcal{D}_{\gamma^*}$ indexed by the free semigroup F_d^+ , so what we mean by the canonical row shift on this space is just moving elements between these copies. Further, to explain the action of V on $\mathfrak{g}$, note that $V^\mathcal{G}$ maps $\mathfrak{g}$ into $\mathfrak{g} \oplus \mathcal{E}^\mathfrak{g}$ and now $\mathbb{1}_{\mathfrak{g}}$ acts as identity on $\mathfrak{g}$, the contraction γ^* maps $\mathcal{E}^\mathfrak{g}$ into $\mathcal{E}_*^\mathfrak{k} \subset \mathcal{K}$ and D_{γ^*} maps $\mathcal{E}^\mathfrak{g}$ into $\mathcal{D}_{\gamma^*}$ which we interpret as $(\mathcal{D}_{\gamma^*})_0 \subset \bigoplus_{\alpha \in F_d^+} (\mathcal{D}_{\gamma^*})_\alpha$.

Note that for $\gamma = 0$ the γ -cascade is nothing but the direct sum of the two processes $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g})$ and $(\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$.

Theorem 4.3. Given processes $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ and $(\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$ (with a common multiplicity space $\mathcal{P}$) and a contraction $\gamma : \mathcal{E}_*^{\mathfrak{k}} \rightarrow \mathcal{E}^{\mathfrak{g}}$. Then the γ -cascade

$$(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \triangleleft_{\gamma} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$$

is a process,

$$\begin{aligned} \gamma &= P_{\mathcal{E}^{\mathfrak{g}}} |_{\mathcal{E}_*^{\mathfrak{k}}} = P_{\mathcal{G}} |_{\mathcal{E}_*^{\mathfrak{k}}} \\ \gamma^* &= P_{\mathcal{E}_*^{\mathfrak{k}}} |_{\mathcal{E}^{\mathfrak{g}}} = P_{\mathcal{K}} |_{\mathcal{E}^{\mathfrak{g}}} \end{aligned}$$

and we have a short exact sequence

$$0 \rightarrow (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \xrightarrow{1_{\mathfrak{g}}} (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \triangleleft_{\gamma} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) \xrightarrow{P_{\mathfrak{k}}} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) \rightarrow 0.$$

In other words, $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ is a subprocess and $(\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$ is a quotient process of the γ -cascade. Conversely, in a short exact sequence

$$0 \longrightarrow (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \xrightarrow{1'_{\mathfrak{g}}} (\mathcal{H}', V', \mathfrak{h}') \xrightarrow{P'_{\mathfrak{k}}} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) \longrightarrow 0,$$

where $1'_{\mathfrak{g}}$ resp. $P'_{\mathfrak{k}}$ are an isometric resp. a coisometric morphism, the process $(\mathcal{H}', V', \mathfrak{h}')$ is isomorphic to a γ -cascade for a contraction $\gamma : \mathcal{E}_*^{\mathfrak{k}} \rightarrow \mathcal{E}^{\mathfrak{g}}$ such that moreover the corresponding extensions are equivalent, i.e. the following diagram commutes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) & \xrightarrow{1'_{\mathfrak{g}}} & (\mathcal{H}', V', \mathfrak{h}') & \xrightarrow{P'_{\mathfrak{k}}} & (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) \longrightarrow 0 \\ & & \uparrow = & & \uparrow j \text{ isom.} & & \uparrow = \\ 0 & \longrightarrow & (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) & \xrightarrow{1_{\mathfrak{g}}} & (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \triangleleft_{\gamma} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) & \xrightarrow{P_{\mathfrak{k}}} & (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) \longrightarrow 0 \end{array}$$

Two cascades with $\gamma_1, \gamma_2 : \mathcal{E}_*^{\mathfrak{k}} \rightarrow \mathcal{E}^{\mathfrak{g}}$ are equivalent as extensions if and only if γ_1 and γ_2 are equal.

We can summarize the theorem by saying that there is a one-to-one correspondence between equivalence classes of extensions of the process $(\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$ by the process $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ and contractions from $\mathcal{E}_*^{\mathfrak{k}}$ to $\mathcal{E}^{\mathfrak{g}}$ and this correspondence is given by the γ -cascade construction.

Proof. Working within the γ -cascade as defined above we have $\mathcal{E}^{\mathfrak{g}}$ identified via the isometry $\begin{pmatrix} \gamma^* \\ D_{\gamma^*} \end{pmatrix}$ with a subspace of $\mathcal{E}_*^{\mathfrak{k}} \oplus (\mathcal{D}_{\gamma^*})_0$. Note that $\mathcal{E}_*^{\mathfrak{k}}$ is the wandering subspace arising from the Wold decomposition of $V^{\mathcal{K}}$, see [33], and $(\mathcal{D}_{\gamma^*})_0$ is another wandering subspace with all translates orthogonal to the translates of $\mathcal{E}_*^{\mathfrak{k}}$. Hence the embedded $\mathcal{E}^{\mathfrak{g}}$ is wandering for V . With this it is now easy to check that V is a row isometry, that $(\mathcal{H}, V, \mathfrak{h})$ is a process with subprocess $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ and quotient process $(\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$ and that γ and γ^* satisfy the formulas stated.

Now suppose that the process $(\mathcal{H}', V', \mathfrak{h}')$ is given by a short exact sequence, i.e. as an extension of $(\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})$ by $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$. We define $\gamma := P_{\mathcal{E}^{\mathfrak{g}}} |_{\mathcal{E}_*^{\mathfrak{k}}} : \mathcal{E}_*^{\mathfrak{k}} \rightarrow \mathcal{E}^{\mathfrak{g}}$ and then form the corresponding γ -cascade $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \triangleleft_{\gamma} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) = (\mathcal{H}, V, \mathfrak{h})$. We verify that this yields an equivalent extension by constructing a unitary $J^* : \mathcal{H}' \rightarrow \mathcal{H}$ which intertwines the row isometries V' and V and which maps $1'_{\mathfrak{g}} \xi$ to $1_{\mathfrak{g}} \xi$ if $\xi \in \mathfrak{g}$ and $(P'_{\mathfrak{k}})^* \eta$ to $(P_{\mathfrak{k}})^* \eta$ if $\eta \in \mathcal{K}$. In fact, then the adjoint $J : \mathcal{H} \rightarrow \mathcal{H}'$ is an extended morphism and its restriction $j : \mathfrak{h} \rightarrow \mathfrak{h}'$ is the isomorphism we look for.

To see that J^* exists it is enough to check that the remaining parts can be matched correctly. We invoke the following lemma which is a standard tool in operator theory.

Lemma 4.4. Let $\mathcal{L}'$ and $\mathcal{L}$ be Hilbert spaces and $\mathcal{L}_0$ a closed subspace of $\mathcal{L}$. If $w : \mathcal{L}' \rightarrow \mathcal{L}$ is an isometry such that $\mathcal{L} = \overline{\text{span}}\{\mathcal{L}_0, w\mathcal{L}'\}$ then w is unitarily equivalent to

$$\begin{pmatrix} \rho \\ D_{\rho} \end{pmatrix} : \mathcal{L}' \rightarrow \mathcal{L}_0 \oplus \mathcal{D}_{\rho} \quad \text{where } \rho = P_{\mathcal{L}_0} w.$$

The unitary from $\mathcal{L}$ to $\mathcal{L}_0 \oplus \mathcal{D}_\rho$ needed in the lemma is the identity on $\mathcal{L}_0$ and it is $w\xi \mapsto \begin{pmatrix} \rho\xi \\ D_\rho\xi \end{pmatrix}$ for $\xi \in \mathcal{L}'$. We apply Lemma 4.4 with $\mathcal{L}' = \mathcal{E}^\mathfrak{g}$, with w being the isometric embedding of $\mathcal{E}^\mathfrak{g}$ into $\mathcal{H}'$ and with $\mathcal{L} = \overline{\text{span}}\{\mathcal{K}, w\mathcal{E}^\mathfrak{g}\}$, $\mathcal{L}_0 = \mathcal{K}$. Lemma 4.1 shows that $\rho = \gamma^*$ in this case and moreover that the embeddings of translates $V'_\alpha \mathcal{E}^\mathfrak{g}$ follow exactly the pattern exposed by the γ -cascade. Hence we can put all the pieces together and get J^* .

Finally, if $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) \triangleleft_{\gamma_1} (\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$ and $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) \triangleleft_{\gamma_2} (\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$ are equivalent extensions then the unitary intertwiner J^* constructed above maps for each element of $\mathcal{E}^\mathfrak{g}$ its first embedding to the second. The same happens to elements of $\mathcal{E}_*^\mathfrak{k}$. Hence if we suppress the embeddings we find that $P_{\mathcal{E}^\mathfrak{g}}|_{\mathcal{E}_*^\mathfrak{k}}$ is the same operator in both cases, i.e., $\gamma_1 = \gamma_2$. $\square$


Remark: It is instructive to look at the situation described in the previous theorem from the point of view of dilation theory. Then we start with a row contraction on $\mathfrak{h} = \mathfrak{g} \oplus \mathfrak{k}$ of the form

$$\begin{pmatrix} X & 0 \\ Y & Z \end{pmatrix}$$

which is called a lifting (in our application this is $p_{\mathfrak{h}}V|_{\mathfrak{h} \otimes \mathcal{P}}$). It is well known that in such a situation Y must have the form $(D_{Z^*})^* \gamma^* D_X$ with a contraction $\gamma : \mathcal{D}_{Z^*} \rightarrow \mathcal{D}_X$, see [18]. Chapter IV, Lemma 2.1 for $d = 1$ and [16], Prop. 3.1 for the general case. Hence a γ -cascade can also be thought of as a description of the structure of the minimal isometric dilation of such a row contraction. Liftings and their dilations are studied in [15, 16] and the results can be interpreted in the language of processes which we use in this paper.

5. CASCADES OF SYSTEMS

One of the things one can do with linear systems is to stick them together in various ways. The most basic way to do that is to take the output of one system I and to use it as the input of another system II . The combined system is then called a *simple cascade*, see for example [19] for the classical theory ($d = 1$). It also works for the noncommutative Fornasini-Marchesini systems we have been considering here. We need a slight generalization where the input of system II is obtained from the output of system I by applying a transformation Γ to it. We call this a Γ -cascade of systems.



We assume here that the two noncommutative Fornasini-Marchesini systems both have the same multiplicity d . Then the internal space of the combined system is defined to be the direct sum of the internal spaces of systems I and II and, with $u_{II}(\alpha) = \Gamma y_I(\alpha)$ (for all $\alpha \in F_d^+$), it follows, by eliminating variables, that the structure maps (A, B, C, D) of the combined system are obtained from the structure maps (A^I, B^I, C^I, D^I) of system I and $(A^{II}, B^{II}, C^{II}, D^{II})$ of system II by

$$A_j = \begin{pmatrix} A_j^I & 0 \\ B_j^{II} \Gamma C^I & A_j^{II} \end{pmatrix}, \quad B_j = \begin{pmatrix} B_j^I \\ B_j^{II} \Gamma D^I \end{pmatrix} \quad (j = 1, \dots, d),$$

$$C = \begin{pmatrix} D^{II} \Gamma C^I & C^{II} \end{pmatrix}, \quad D = D^{II} \Gamma D^I.$$

In this case we also speak of a Γ -cascade of structure maps.

The transfer function $\mathcal{T}$ of such a Γ -cascade of systems (or of structure maps) factorizes. If T is a power series with coefficients $T_\alpha \in \mathcal{B}(V, W)$ and Γ maps W to W' then we denote by ΓT the power series with coefficients $\Gamma T_\alpha \in \mathcal{B}(V, W')$. With this convention it is not difficult to check that the transfer function $\mathcal{T}$ of the combined system is obtained from the transfer functions $\mathcal{T}^{II}$ and $\mathcal{T}^I$ of systems II and I by

$$\mathcal{T}(z) = \mathcal{T}^{II}(z) \Gamma \mathcal{T}^I(z)$$

(with $z^\alpha z^\beta = z^{\beta\alpha}$).

Now we prove that if $(\mathcal{H}, V, \mathfrak{h})$ is a γ -cascade $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) \triangleleft_\gamma (\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$ of processes then we can think of its Fornasini-Marchesini system, from Proposition 3.2, as a cascade of the systems associated to $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g})$ and $(\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$. This is not so obvious if you arrive at the notion of a γ -cascade of processes from a dilation point of view but it nicely clarifies the system theory involved and of course it motivates our choice of terminology.

We assume that we have structure maps $(A^\mathcal{G}, B^\mathcal{G}, C^\mathcal{G}, D^\mathcal{G})$ represented by the process $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g})$ and $(A^\mathcal{K}, B^\mathcal{K}, C^\mathcal{K}, D^\mathcal{K})$ represented by the process $(\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$, according to Definition 3.1.

Now we form a γ -cascade of processes

$$(\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) \triangleleft_\gamma (\mathcal{K}, V^\mathcal{K}, \mathfrak{k}) = (\mathcal{H}, V, \mathfrak{h})$$

and we introduce output and input spaces represented by $(\mathcal{H}, V, \mathfrak{h})$:

$$\mathcal{Y}_0 := \mathcal{Y}_0^\mathcal{G}, \quad \mathcal{U}_0 := \mathcal{U}_0^\mathcal{K}.$$

We have the inclusions $\mathcal{U}_0^\mathcal{K} \subset \mathcal{E}^\mathfrak{k} \subset \mathcal{E}$ and $\mathcal{Y}_0^\mathcal{G} \subset \mathfrak{g} \oplus \mathcal{E}^\mathfrak{g} \subset \mathfrak{h} \oplus \mathcal{E}$ and $\mathcal{Y}_0^\mathcal{G}$ is wandering for $V^\mathcal{G}$ and hence for V , so indeed this yields a representation of structure maps (A, B, C, D) by the process $(\mathcal{H}, V, \mathfrak{h})$. Further we define

$$\Gamma := P_{\mathcal{U}_0^\mathcal{G}}|_{\mathcal{Y}_0^\mathcal{K}} = P_{\mathcal{U}_0^\mathcal{G}}\gamma P_{\mathcal{E}_*^\mathfrak{k}}|_{\mathcal{Y}_0^\mathcal{K}}$$

where the latter equality follows from $\mathcal{U}_0^\mathcal{G} \subset \mathcal{E}^\mathfrak{g}$ together with the geometry of the γ -cascade (see Definition 4.2 and Lemma 4.1).

A subspace $\mathcal{L}$ of $\mathcal{H}$ is called a *left support* of γ if $P_\mathcal{L}\gamma P_{\mathcal{E}_*^\mathfrak{k}} = \gamma P_{\mathcal{E}_*^\mathfrak{k}}$ and it is called a *right support* of γ if $\gamma P_{\mathcal{E}_*^\mathfrak{k}} = \gamma P_{\mathcal{E}_*^\mathfrak{k}} P_\mathcal{L}$. With these preparations we can now find a Γ -cascade of systems in the γ -cascade of processes.

Theorem 5.1. Suppose that $\mathcal{U}_0^\mathcal{G}$ is a left support and $\mathcal{Y}_0^\mathcal{K}$ is a right support of γ . Then the structure maps (A, B, C, D) are a Γ -cascade of $(A^\mathcal{K}, B^\mathcal{K}, C^\mathcal{K}, D^\mathcal{K})$ and $(A^\mathcal{G}, B^\mathcal{G}, C^\mathcal{G}, D^\mathcal{G})$. Explicitly (for $j = 1, \dots, d$):

$$\begin{aligned} A_j &= \begin{pmatrix} A_j^\mathcal{K} & 0 \\ B_j^\mathcal{G} \Gamma C^\mathcal{K} & A_j^\mathcal{G} \end{pmatrix} & B_j &= \begin{pmatrix} B_j^\mathcal{K} \\ B_j^\mathcal{G} \Gamma D^\mathcal{K} \end{pmatrix} \\ C &= (D^\mathcal{G} \Gamma C^\mathcal{K} \quad C^\mathcal{G}), & D &= D^\mathcal{G} \Gamma D^\mathcal{K}. \end{aligned}$$

Roughly speaking, $\mathcal{U}_0^\mathcal{G}$ left and $\mathcal{Y}_0^\mathcal{K}$ right supporting γ means that they are chosen big enough to transport the information contained in γ . An example is given by the choice $\mathcal{U}_0^\mathcal{G} := \mathcal{E}^\mathfrak{g}$ and $\mathcal{Y}_0^\mathcal{K} := \mathcal{E}_*^\mathfrak{k}$ and in this case $\Gamma = \gamma$.

Proof. We verify the explicit formulas step by step. The arguments are based on the geometry of a γ -cascade as given in Definition 4.2, in particular: $V|_\mathfrak{g} = V^\mathcal{G}|_\mathfrak{g}$ maps $\mathfrak{g}$ into $\mathfrak{g} \oplus \mathcal{E}^\mathfrak{g}$ and $(\mathfrak{g} \oplus \mathcal{E}^\mathfrak{g}) \ominus V^\mathcal{G}(\mathfrak{g} \otimes \mathcal{P})$ is orthogonal to the range of $V^\mathcal{G}$, further $P_{\mathfrak{g} \oplus \mathcal{E}^\mathfrak{g}}|_\mathcal{K} = P_{\mathcal{E}^\mathfrak{g}}|_\mathcal{K} = \gamma P_{\mathcal{E}_*^\mathfrak{k}}|_\mathcal{K}$.

$$\begin{aligned} P_\mathfrak{k} A_j &= P_\mathfrak{k} V_j^*|_\mathfrak{h} = (P_\mathfrak{k} V_j^*|_\mathfrak{k} \quad P_\mathfrak{k} V_j^*|_\mathfrak{g}) = (A_j^\mathcal{K} \quad 0) \\ P_\mathfrak{g} A_j &= P_\mathfrak{g} V_j^*|_\mathfrak{h} = (P_\mathfrak{g} V_j^*|_\mathfrak{k} \quad P_\mathfrak{g} V_j^*|_\mathfrak{g}) = (V_j^{\mathcal{G}*} P_{\mathfrak{g} \oplus \mathcal{E}^\mathfrak{g}}|_\mathfrak{k} \quad V_j^{\mathcal{G}*}|_\mathfrak{g}) \\ &= (V_j^{\mathcal{G}*} P_{\mathcal{E}^\mathfrak{g}} \gamma P_{\mathcal{E}_*^\mathfrak{k}}|_\mathfrak{k} \quad V_j^{\mathcal{G}*}|_\mathfrak{g}) = (V_j^{\mathcal{G}*} P_{\mathcal{U}_0^\mathcal{G}} P_{\mathcal{E}^\mathfrak{g}} \gamma P_{\mathcal{E}_*^\mathfrak{k}} P_{\mathcal{Y}_0^\mathcal{K}}|_\mathfrak{k} \quad V_j^{\mathcal{G}*}|_\mathfrak{g}) \\ &= (B_j^\mathcal{G} \Gamma C^\mathcal{K} \quad A_j^\mathcal{G}) \\ P_\mathfrak{k} B_j &= P_\mathfrak{k} V_j^*|_{\mathcal{U}_0^\mathcal{K}} = P_\mathfrak{k} V_j^{\mathcal{K}*}|_{\mathcal{U}_0^\mathcal{K}} = B_j^\mathcal{K} \\ P_\mathfrak{g} B_j &= P_\mathfrak{g} V_j^*|_{\mathcal{U}_0^\mathcal{K}} = P_\mathfrak{g} V_j^{\mathcal{G}*} P_{\mathfrak{g} \oplus \mathcal{E}^\mathfrak{g}}|_{\mathcal{U}_0^\mathcal{K}} = P_\mathfrak{g} V_j^{\mathcal{G}*} P_{\mathcal{U}_0^\mathcal{G}} P_{\mathcal{E}^\mathfrak{g}} \gamma P_{\mathcal{E}_*^\mathfrak{k}} P_{\mathcal{Y}_0^\mathcal{K}}|_{\mathcal{U}_0^\mathcal{K}} = B_j^\mathcal{G} \Gamma D^\mathcal{K} \\ C &= (P_{\mathcal{Y}_0^\mathcal{G}}|_\mathfrak{k} \quad P_{\mathcal{Y}_0^\mathcal{G}}|_\mathfrak{g}) = (P_{\mathcal{Y}_0^\mathcal{G}} P_{\mathcal{U}_0^\mathcal{G}} P_{\mathcal{E}^\mathfrak{g}} \gamma P_{\mathcal{E}_*^\mathfrak{k}} P_{\mathcal{Y}_0^\mathcal{K}}|_\mathfrak{k} \quad P_{\mathcal{Y}_0^\mathcal{G}}|_\mathfrak{g}) = (D^\mathcal{G} \Gamma C^\mathcal{K} \quad C^\mathcal{G}) \\ D &= P_{\mathcal{Y}_0^\mathcal{K}}|_{\mathcal{U}_0^\mathcal{K}} = P_{\mathcal{Y}_0^\mathcal{K}}|_{\mathcal{U}_0^\mathcal{K}} = P_{\mathcal{Y}_0^\mathcal{K}} P_{\mathcal{U}_0^\mathcal{G}} P_{\mathcal{E}^\mathfrak{g}} \gamma P_{\mathcal{E}_*^\mathfrak{k}} P_{\mathcal{Y}_0^\mathcal{K}}|_{\mathcal{U}_0^\mathcal{K}} = D^\mathcal{G} \Gamma D^\mathcal{K}. \end{aligned}$$

□

It is possible to iterate the construction shown in Theorem 5.1. Because $V^* \mathcal{E}_*^{\mathfrak{g}} = V^{\mathcal{G}*} \mathcal{E}_*^{\mathfrak{g}} = \{0\}$ we have $\mathcal{E}_*^{\mathfrak{g}} \subset \mathcal{E}_*$. Hence if $(\mathcal{F}, V^{\mathcal{F}}, \mathfrak{f})$ is another process (with the same multiplicity) and $\gamma_2: \mathcal{E}_*^{\mathfrak{g}} \rightarrow \mathcal{E}^{\mathfrak{f}}$ is a contraction then not only can we form $(\mathcal{F}, V^{\mathcal{F}}, \mathfrak{f}) \triangleleft_{\gamma_2} (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ but also $(\mathcal{F}, V^{\mathcal{F}}, \mathfrak{f}) \triangleleft_{\gamma_2} [(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \triangleleft_{\gamma} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k})]$ with $\hat{\gamma}_2: \mathcal{E}_* \rightarrow \mathcal{E}^{\mathfrak{f}}$ given by

$$\hat{\gamma}_2 := \begin{cases} \gamma_2 & \text{on } \mathcal{E}_*^{\mathfrak{g}} \\ 0 & \text{on } \mathcal{E}_* \ominus \mathcal{E}_*^{\mathfrak{g}} \end{cases}$$

Theorem 5.1 applies iteratively. This may be seen as another justification for calling such a structure a cascade.

6. OBSERVABILITY AND ASYMPTOTIC COMPLETENESS

To make use of the system theory now available to us for the study of processes we discuss the control theory concept of observability in the multi-variable setting. See for example [2] for a recent treatment in a purely operator theoretic spirit.

Definition 6.1. Given an output pair (A, C) for an internal space $\mathcal{X}$ and an output space $\mathcal{Y}$, a subset $\mathcal{X}' \subset \mathcal{X}$ is called observable if $(CA^\alpha|_{\mathcal{X}'})_{\alpha \in F_d^+}$, the observability map restricted to $\mathcal{X}'$, is injective (as a map from $\mathcal{X}'$ to the $\mathcal{Y}$ -valued functions on F_d^+).

If (A, C) is represented by the process $(\mathcal{H}, V, \mathfrak{h})$ then we also say in this case that $\mathcal{X}' (\subset \mathfrak{h})$ is observable in $(\mathcal{H}, V, \mathfrak{h})$ by the output space $\mathcal{Y}_0$.

The interpretation of observability is that every $\xi \in \mathcal{X}'$ can be reconstructed from the outputs $CA^\alpha \xi$.

Proposition 6.2. The subset $\mathcal{X}' \subset \mathfrak{h}$ is observable in $(\mathcal{H}, V, \mathfrak{h})$ by the output space $\mathcal{Y}_0$ if and only if $P_{\mathcal{Y}_+}|_{\mathcal{X}'}$ is injective.

Proof. For $\xi \in \mathcal{X}'$ we have

$$CA^\alpha \xi = P_{\mathcal{Y}_0} V_\alpha^* \xi = V_\alpha^* P_{\mathcal{Y}_\alpha} \xi$$

and we conclude that $\mathcal{X}'$ is observable if and only if $(P_{\mathcal{Y}_\alpha}|_{\mathcal{X}'})_{\alpha \in F_d^+}$ is injective. Because the projection $P_{\mathcal{Y}_+}$ is the supremum of the projections $(P_{\mathcal{Y}_\alpha})_{\alpha \in F_d^+}$ we can replace the family $(P_{\mathcal{Y}_\alpha}|_{\mathcal{X}'})_{\alpha \in F_d^+}$ by the single contraction $P_{\mathcal{Y}_+}|_{\mathcal{X}'}$. □

If we have a γ -cascade and we use the maximal input space $\mathcal{E}^{\mathfrak{g}}$ of the subprocess as output space for the cascade, roughly speaking if we confine our observations to the subprocess, then questions of observability become an interesting issue of theoretical and practical importance.

Theorem 6.3. Consider the γ -cascade

$$(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \triangleleft_{\gamma} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) = (\mathcal{H}, V, \mathfrak{h})$$

and the output space $\mathcal{Y}_0 := \mathcal{E}^{\mathfrak{g}}$. The following assertions are equivalent:

- (1a) $\mathfrak{k}$ is observable in $(\mathcal{H}, V, \mathfrak{h})$ by $\mathcal{Y}_0$.
- (1b) $P_{\mathcal{Y}_+}|_{\mathfrak{k}}$ is injective.
- (2a) $\overline{\text{span}}\{(A^\alpha)^* \mathfrak{g}: \alpha \in F_d^+\} = \mathfrak{h}$
- (2b) $P_{\mathfrak{h}}|_{\mathcal{G}}: \mathcal{G} \rightarrow \mathfrak{h}$ has dense range.
- (3a) $\mathcal{G} = \mathcal{H}$
- (3b) There is an extended morphism associated to the morphism $\mathbb{1}_{\mathfrak{g}}: \mathfrak{g} \rightarrow \mathfrak{h}$ which is a unitary from $\mathcal{H}$ to $\mathcal{H}$.
- (4a) $V^{\mathcal{K}}$ is a row shift and $\gamma: \mathcal{E}_*^{\mathfrak{k}} \rightarrow \mathcal{E}^{\mathfrak{g}}$ is injective.
- (4b) $V^{\mathcal{K}}$ is a row shift and $\gamma: \mathcal{E}_*^{\mathfrak{k}} \rightarrow \mathcal{E}^{\mathfrak{g}}$ is isometric.

If the transition operator Z of $(\mathcal{H}, V, \mathfrak{h})$ is unital then we also have the following equivalent condition:

- (5) $\lim_{n \rightarrow \infty} Z^n(P_{\mathfrak{g}}) = \mathbb{1}_{\mathfrak{h}}$ (in the strong operator topology)

Proof. The equivalence of (1a) and (1b) is Proposition 6.2. Further

$$\begin{aligned} & P_{\mathcal{Y}_+}|_{\mathfrak{k}} \text{ injective} \\ \Leftrightarrow & P_{\mathcal{G}}|_{\mathfrak{h}} \text{ injective} \quad (\text{by adding the space } \mathfrak{g}) \\ \Leftrightarrow & P_{\mathfrak{h}}|_{\mathcal{G}} \text{ has dense range} \quad (\text{adjoint map}) \end{aligned}$$

which shows (1b) $\Leftrightarrow$ (2b). Because $G = \overline{\text{span}}\{V_\alpha \mathfrak{g} : \alpha \in F_d^+\}$ assertion (2b) is equivalent to

$$\mathfrak{h} = \overline{\text{span}}\{P_{\mathfrak{h}} V_\alpha \mathfrak{g} : \alpha \in F_d^+\} = \overline{\text{span}}\{(A^\alpha)^* \mathfrak{g} : \alpha \in F_d^+\}$$

which is (2a). On the other hand

$$P_{\mathfrak{h}}|_{\mathcal{G}} \text{ has dense range} \Rightarrow P_{\mathcal{H}}|_{\mathcal{G}} \text{ has dense range}$$

because $\mathcal{H}$ is the closed linear span of the $V_\alpha \mathfrak{h}$ with $\alpha \in F_d^+$ and we have $\mathcal{G} \supset V_\alpha \mathcal{G}$, hence

$$P_{V_\alpha \mathfrak{h}} \mathcal{G} \supset P_{V_\alpha \mathfrak{h}} V_\alpha \mathcal{G} = V_\alpha P_{\mathfrak{h}} \mathcal{G}$$

which is dense in $V_\alpha \mathfrak{h}$. But $\mathcal{G}$ is a closed subspace of $\mathcal{H}$ and we conclude that (2b) implies (3a). The converse, (3a) implies (2b), is obvious.

It is easy to check that in a γ -cascade the morphism $\mathbb{1}_{\mathfrak{g}}$, the embedding of $\mathfrak{g}$ into $\mathfrak{h}$, always has $\mathbb{1}_{\mathcal{G}}$, the embedding of $\mathcal{G}$ into $\mathcal{H}$, as an associated extended morphism. If we have (3a), i.e. $\mathcal{G} = \mathcal{H}$, then $\mathbb{1}_{\mathcal{G}}$ is nothing but the identity operator $\mathbb{1}_{\mathcal{H}}$ on $\mathcal{H}$, hence (3a) implies (3b). Conversely (3a) is implicit in the statement of (3b).

From Definition 4.2 of the γ -cascade we have

$$\mathcal{E}^{\mathfrak{g}} \subset \mathcal{E}_*^{\mathfrak{k}} \oplus (D_{\gamma^*})_0$$

with $\mathcal{E}_*^{\mathfrak{k}} \subset \mathcal{K}$ and $(D_{\gamma^*})_0 \subset \mathcal{K}^\perp$, from which we get $P_{\mathcal{K}} \mathcal{Y}_+ \subset \bigoplus_{\alpha} V_\alpha \mathcal{E}_*^{\mathfrak{k}}$. If we have (3a), i.e. $\mathcal{G} = \mathcal{H}$, then, because $\mathcal{G} = \mathfrak{g} \oplus \mathcal{Y}_+$ and $\mathfrak{g} \perp \mathcal{K}$, we find $P_{\mathcal{K}} \mathcal{Y}_+ = \mathcal{K}$ and hence $\mathcal{K} = \bigoplus_{\alpha} V_\alpha \mathcal{E}_*^{\mathfrak{k}}$ which means that $V^{\mathcal{K}}$ is a row shift. From (3a) we have $\mathcal{K} \subset \mathcal{G}$ and together with $\mathcal{E}_*^{\mathfrak{k}} \perp \mathfrak{g}$ and $\mathcal{E}_*^{\mathfrak{k}} \perp V_\alpha \mathcal{E}^{\mathfrak{g}}$ for all $\alpha \neq 0$ (by Lemma 4.1) we conclude that $\mathcal{E}_*^{\mathfrak{k}}$ is actually a subspace of $\mathcal{E}^{\mathfrak{g}}$ which means that $\gamma = P_{\mathcal{E}^{\mathfrak{g}}}|_{\mathcal{E}_*^{\mathfrak{k}}}$ is an isometric embedding. Hence (3a) implies (4b). Obviously (4b) implies (4a) and from (4a), together with Lemma 4.1, we get an injective map

$$P_{\mathcal{Y}_+}|_{\mathcal{K}} = \bigoplus_{\alpha} V_\alpha \gamma V_\alpha^* : \quad \mathcal{K} = \bigoplus_{\alpha} V_\alpha \mathcal{E}_*^{\mathfrak{k}} \rightarrow \bigoplus_{\alpha} V_\alpha \mathcal{E}^{\mathfrak{g}} = \mathcal{Y}_+$$

and (1b) follows.

Finally if Z is unital then the projections $\theta^n(P_{\mathfrak{g}})$ increase with n and converge to their supremum $P_{\mathcal{G}}$ (see Section 2), hence $Z^n(P_{\mathfrak{g}}) = P_{\mathfrak{h}} \theta^n(P_{\mathfrak{g}})|_{\mathfrak{h}}$ converges to $P_{\mathfrak{h}} P_{\mathcal{G}}|_{\mathfrak{h}}$ (in the strong operator topology). Now (3a) implies that $P_{\mathfrak{h}} P_{\mathcal{G}}|_{\mathfrak{h}} = \mathbb{1}_{\mathfrak{h}}$ and this implies (2b), hence all these assertions, including (5), are equivalent. $\square$

Remark: Of course (3b) does not mean that $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ and $(\mathcal{H}, V, \mathfrak{h})$ are isomorphic as processes. In fact, if $\mathfrak{g} \neq \mathfrak{h}$ then the morphism $\mathbb{1}_{\mathfrak{g}}$, the embedding of $\mathfrak{g}$ into $\mathfrak{h}$, is not unitary.

Definition 6.4. The short exact sequence

$$0 \longrightarrow (\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g}) \xrightarrow{\mathbb{1}_{\mathfrak{g}}} (\mathcal{H}, V, \mathfrak{h}) \xrightarrow{P_{\mathfrak{k}}} (\mathcal{K}, V^{\mathcal{K}}, \mathfrak{k}) \longrightarrow 0$$

is called *asymptotically complete* if one (and hence all) of the assertions in Theorem 6.3 are satisfied.

This terminology is borrowed from scattering theory. It will be explained in the final section of this paper how the setting above is related to already existing scattering theories for Markov chains.

Corollary 6.5. Let an asymptotically complete sequence as in Definition 6.4 be given together with an input space $\mathcal{U}_0$ and the output space $\mathcal{Y}_0 = \mathcal{E}^{\mathfrak{g}}$, both represented by $(\mathcal{H}, V, \mathfrak{h})$. Then $\mathcal{O}_{C,A}|_{\mathfrak{k}} = (CA^\alpha|_{\mathfrak{k}})_{\alpha \in F_d^+} = (C(A^{\mathcal{K}})^\alpha)_{\alpha \in F_d^+}$, the restriction of the observability map to $\mathfrak{k}$, is isometric and the transfer function $\mathcal{T}$ is inner, i.e., the corresponding multi-analytic operator is isometric.

Proof. Note that $\mathcal{Y}_0 = \mathcal{E}^{\mathfrak{g}} \perp \mathfrak{g}$ indeed gives us $CA^\alpha|_{\mathfrak{k}} = C(A^\mathcal{K})^\alpha$ for all α . We have $V_\alpha CA^\alpha|_{\mathfrak{k}} = P_{\mathcal{Y}_\alpha}|_{\mathfrak{k}}$. The multi-analytic operator corresponding to the transfer function $\mathcal{T}$ is $P_{\mathcal{Y}_+}|_{\mathcal{U}_+}$.

From asymptotic completeness it follows that $\mathfrak{g} \oplus \mathcal{Y}_+ = \mathcal{G} = \mathcal{H}$, see Theorem 6.3 (3a). Hence, because $\mathfrak{k}$ and $\mathcal{U}_+$ are orthogonal to $\mathfrak{g}$, they are both subspaces of $\mathcal{Y}_+$ and it follows that $P_{\mathcal{Y}_+}|_{\mathfrak{k}}$ and $P_{\mathcal{Y}_+}|_{\mathcal{U}_+}$ are isometric. $\square$

We describe a few alternative ways to interpret asymptotic completeness. First, if we are in an asymptotically complete situation and we choose $\mathcal{U}_0$ maximal, i.e. $\mathcal{U}_0 = \mathcal{E}$, then the sum of the identity $\mathbb{1}_{\mathfrak{g}}$ (on $\mathfrak{g}$), of the observability map $\mathcal{O}_{C,A}|_{\mathfrak{k}}$ (on $\mathfrak{k}$) and of the multi-analytic operator associated to $\mathcal{T}$ (on $\mathcal{U}_+$) is equal to the identity $\mathbb{1}_{\mathcal{H}}$. This is a direct consequence of Corollary 6.5. But it is the identity $\mathbb{1}_{\mathcal{H}}$ presented with a change of basis that describes the relative position of the weak filtrations of the process $(\mathcal{H}, V, \mathfrak{h})$ and of the process $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$.

Second, from the point of view of dilation theory, in the case of asymptotic completeness we deal with so-called subisometric dilations. We don't go into this here, see [16] for details.

Third, asymptotic completeness and its observability aspect can also be given an explicit quantum physical interpretation by reconsidering the observables discussed in Proposition 2.6. Indeed, in the case of asymptotic completeness the Hilbert space $\mathcal{H}$ is the closed span of the spaces $V^{(n)} \mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}$ (with $n \in \mathbb{N}_0$, the case $n = 0$ to be interpreted as $\mathfrak{g}$). It follows that the algebras $V^{(n)} \mathcal{B}(\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}) V^{(n)*}$ generate $\mathcal{B}(\mathcal{H})$ (as a von Neumann algebra). On the other hand, the observables

$$\begin{aligned} X_n &= V^{(n)} X \otimes \bigotimes_1^n \mathbb{1}_{\mathcal{P}} V^{(n)*}, \\ Y_{m,n} &= V^{(n)} \mathbb{1}_{\mathcal{H}} \otimes \bigotimes_1^{m-1} \mathbb{1}_{\mathcal{P}} \otimes Y \otimes \bigotimes_{m+1}^n \mathbb{1}_{\mathcal{P}} V^{(n)*}, \quad m = 1, \dots, n, \end{aligned}$$

if we consider all $X \in \mathcal{B}(\mathfrak{g}) = P_{\mathfrak{g}} \mathcal{B}(\mathfrak{h}) P_{\mathfrak{g}}$ and all $Y \in \mathcal{B}(\mathcal{P})$, generate $V^{(n)} \mathcal{B}(\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}) V^{(n)*}$ (as a von Neumann algebra). Conclusion: We can approximate arbitrary observables in $\mathcal{B}(\mathcal{H})$ in the weak (or strong) operator topology by observables generated by these X_n and $Y_{m,n}$. In particular, in physics language, we can answer all questions about observables on the part $\mathfrak{k}$ (and hence also find the state) from measuring observables on the orthogonal part $\mathfrak{g}$ of the internal space (the X_n) together with field observables (the $Y_{m,n}$). We have the following quantitative statement about these approximations which stresses the role of the observability operator $\mathcal{O}_{C,A}$: If $\xi \in \mathfrak{k}$ then the norm distance squared of ξ from the space $\mathfrak{g}_n = \mathfrak{g} \oplus \bigoplus_{|\alpha| < n} V_\alpha \mathcal{E}^{\mathfrak{g}}$ is given by $\|\xi\|^2 - \sum_{|\alpha| < n} \|CA^\alpha \xi\|^2$. If we have asymptotic completeness then $\mathcal{O}_{C,A}|_{\mathfrak{k}}$ is isometric and this distance tends to 0 for $n \rightarrow \infty$.

7. SUBPROCESSES FROM NORMAL INVARIANT STATES

The following way of finding subprocesses gives a connection to a topic which is of natural interest for quantum Markov processes and more general for quantum probability: invariant states. Recall that if ϕ is a normal state of a von Neumann algebra then in this von Neumann algebra there exists a smallest orthogonal projection p such that $\phi(p) = 1$, called the support projection $s(\phi)$. For all elements x we have

$$\phi(x) = \phi(s(\phi)x) = \phi(xs(\phi)) = \phi(s(\phi)xs(\phi)).$$

Essentially the following is Lemma 6.1 from [14], for convenience we include a proof which uses our now familiar terminology and notation.

Proposition 7.1. Given a process $(\mathcal{H}, V, \mathfrak{h})$, suppose that ϕ is a normal state of $\mathcal{B}(\mathfrak{h})$ which is invariant for the transition operator Z , i.e.,

$$\phi(Z(x)) = \phi(x) \quad \text{for all } x \in \mathcal{B}(\mathfrak{h}).$$

Then with $P_{\mathfrak{g}} := s(\phi)$ the subspace $\mathfrak{g}$ is co-invariant for V .

Proof. Let $\mathfrak{k} := \mathfrak{h} \ominus \mathfrak{g}$. Then

$$0 = \phi(P_{\mathfrak{k}}) = \phi(Z(P_{\mathfrak{k}})) = \phi(P_{\mathfrak{g}}Z(P_{\mathfrak{k}})P_{\mathfrak{g}}),$$

hence $P_{\mathfrak{g}}Z(P_{\mathfrak{k}})P_{\mathfrak{g}} = 0$ (because ϕ restricted to the subalgebra $s(\phi)\mathcal{B}(\mathfrak{h})s(\phi)$, obtained by compression with its support, is a faithful state). But

$$P_{\mathfrak{g}}Z(P_{\mathfrak{k}})P_{\mathfrak{g}} = P_{\mathfrak{g}}V(P_{\mathfrak{k}} \otimes \mathbb{1}_{\mathcal{P}})V^*P_{\mathfrak{g}} = X^*X$$

with $X = (P_{\mathfrak{k}} \otimes \mathbb{1}_{\mathcal{P}})V^*P_{\mathfrak{g}}$. Hence $X = 0$ which shows that $\mathfrak{g}$ is co-invariant for V . $\square$

Using Proposition 7.1 we can always find a subprocess from a normal invariant state and this subprocess is nontrivial, in the sense that $\mathfrak{g} \neq \mathfrak{h}$, if and only if the state is not faithful. Definition 6.4 for asymptotic completeness applies and we can use the criteria in Theorem 6.3. Instead of further analyzing the general case we concentrate for the rest of this section on an application to a class of noncommutative Markov processes which are not originally weak Markov processes but which nevertheless can be studied successfully by our methods.

The following construction represents the most basic way of introducing stationary Markov chains in an operator algebraic context. In this form it is taken from [20], see more details there. A more leisurely introduction to the topic is [31]. Let $\mathcal{A}$ and $\mathcal{C}$ be C^* -algebras and let

$$j : \mathcal{A} \rightarrow \mathcal{A} \otimes \mathcal{C}$$

be a non-zero $*$ -homomorphism. This can be iterated to yield the $*$ -homomorphisms

$$j_n : \mathcal{A} \rightarrow \mathcal{A} \otimes \bigotimes_1^n \mathcal{C},$$

where $j_1(a) := j(a) =: \sum_k a_k \otimes c_k \in \mathcal{A} \otimes \mathcal{C}$ and then inductively

$$j_n(a) := \sum_k j_{n-1}(a_k) \otimes c_k \in (\mathcal{A} \otimes \bigotimes_1^{n-1} \mathcal{C}) \otimes \mathcal{C} = \mathcal{A} \otimes \bigotimes_1^n \mathcal{C}.$$

If ϕ resp. ψ are states on $\mathcal{A}$ respectively $\mathcal{C}$ and we impose the stationarity condition

$$(\phi \otimes \psi) \circ j = \phi$$

then we can think of the sequence (j_n) as noncommutative random variables which form a noncommutative stationary Markov chain.

We can obtain an associated weak process in the sense of this paper by applying the GNS-construction which from $(\mathcal{A}, \phi)$ produces $(\mathfrak{h}, \Omega_\phi)$ and from $(\mathcal{C}, \psi)$ produces $(\mathcal{P}, \Omega_\psi)$, the GNS-Hilbert space and a cyclic vector representing the state (in each case). Then the stationarity condition translates into the fact that

$$\begin{aligned} v_1 : \mathfrak{h} &\rightarrow \mathfrak{h} \otimes \mathcal{P} \\ a\Omega_\phi &\mapsto j(a)\Omega_\phi \otimes \Omega_\psi \end{aligned}$$

(with $a \in \mathcal{A}$) is an isometry. In Chapter 1 of [21] v_1 is called the *associated isometry* and it plays a central role there in the analysis of the Markov chain. We are now in a position to deepen this analysis and to make the conceptual framework more elegant by putting it into the context provided in this paper. We start with the adjoint

$$v_1^* : \mathfrak{h} \otimes \mathcal{P} \rightarrow \mathfrak{h}$$

which is a (row) coisometry. Hence its minimal isometric dilation

$$V : \mathcal{H} \otimes \mathcal{P} \rightarrow \mathcal{H}$$

is a row unitary and $(\mathcal{H}, V, \mathfrak{h})$ is a unital process. (It is a well known fact in dilation theory which can be checked directly that the minimal isometric dilation V on $\mathcal{H}$ of a row contraction on $\mathfrak{h}$ is a row unitary if and only if the row contraction is coisometric.) We call $(\mathcal{H}, V, \mathfrak{h})$ the (weak) process

associated to the original stationary Markov chain.

Remark: If we think of V^* as a kind of (Schrödinger) dynamics of vector states for a weak process then we see that for an associated weak process this is provided on $\mathfrak{h}$ by the associated isometry v_1 . Note however that if we consider the original noncommutative random variables j_n as steps of a (Heisenberg) dynamics of observables then there is a time reversal involved if we go to the noncommutative random variables $J^{(n)}$ (as in Section 2) as steps of a (Heisenberg) dynamics of observables for the associated weak process.

But there is more structure available from a stationary Markov chain. One can check that the vector state represented by Ω_ϕ is invariant for the transition operator Z of the (weak) process and hence we can invoke Proposition 7.1 to see that its support, the one-dimensional subspace $\mathfrak{g} = \mathbb{C}\Omega_\phi$, is co-invariant for V . In fact it is easy to check this directly: Consider the one-dimensional subspace $\mathfrak{g} := \mathbb{C}\Omega_\phi \subset \mathfrak{h}$. Because $V^*|_{\mathfrak{h}} = v_1$ and $v_1\Omega_\phi = \Omega_\phi \otimes \Omega_\psi$ we find

$$V^*\mathfrak{g} = V^*\mathbb{C}\Omega_\phi = \mathbb{C}\Omega_\phi \otimes \Omega_\psi \subset \mathfrak{g} \otimes \mathcal{P}$$

which implies that $\mathfrak{g}$ is co-invariant for V and hence we have a subprocess $(\mathcal{G}, V^\mathcal{G}, \mathfrak{g})$. Combined with the analysis in Section 4 we obtain the surprising result that the GNS-Hilbert space of a stationary Markov chain always carries automatically the structure of a γ -cascade

$$(\mathcal{H}, V, \mathfrak{h}) = (\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) \triangleleft_\gamma (\mathcal{K}, V^\mathcal{K}, \mathfrak{k})$$

and we can think of the whole short exact sequence

$$0 \rightarrow (\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) \xrightarrow{1_\mathfrak{g}} (\mathcal{G}, V^\mathcal{G}, \mathfrak{g}) \triangleleft_\gamma (\mathcal{K}, V^\mathcal{K}, \mathfrak{k}) \xrightarrow{P_\mathfrak{k}} (\mathcal{K}, V^\mathcal{K}, \mathfrak{k}) \rightarrow 0.$$

as being *associated* to the stationary Markov chain. We have $\mathfrak{g} = \mathbb{C}\Omega_\phi$ and $\mathfrak{k} = \overline{\mathcal{A}\Omega_\phi} \ominus \mathbb{C}\Omega_\phi$.

Let us investigate what asymptotic completeness means in this case. We can define the associated isometry v_n for the n -th noncommutative random variable j_n which, because it arises from the same iterative procedure, can be expressed by the iteration $V^{(n)}$ of V . With $a \in \mathcal{A}$ and $\xi := a\Omega_\phi \in \mathfrak{h}$ we have

$$j_n(a)\Omega_\phi \otimes \bigotimes_1^n \Omega_\psi = v_n\xi = V^{(n)*}\xi$$

and from that

$$\begin{aligned} P_{\Omega_\phi \otimes \bigotimes_1^n \mathcal{P}} j_n(a)\Omega_\phi \otimes \bigotimes_1^n \Omega_\psi &= P_{\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}} v_n\xi \\ &= V^{(n)*}V^{(n)}P_{\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}} V^{(n)*}\xi = V^{(n)*}P_{V^{(n)}\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}} \xi \end{aligned}$$

which yields the norm equality

$$\|P_{\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}} v_n\xi\| = \|P_{V^{(n)}\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}} \xi\|.$$

Because $P_{V^{(n)}\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}}$ increases to the projection $P_\mathcal{G}$ (which is the limit for $n \rightarrow \infty$ in the strong operator topology) it follows that the property $\mathcal{G} = \mathcal{H}$, equivalent to asymptotic completeness by Theorem 6.3, is also equivalent to

$$\|P_{\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P}} v_n\xi\| \rightarrow \|\xi\| \quad \text{for } n \rightarrow \infty$$

for all $\xi \in \mathfrak{h}$. If the GNS-representation is faithful then we can interpret the Hilbert space norm as a norm $\|\cdot\|_2$ on the C^* -algebra and we can write

$$\|E_{\mathbf{1}_\mathcal{A} \otimes \bigotimes_1^n \mathcal{C}} j_n(a)\|_2 \rightarrow \|a\|_2 \quad \text{for } n \rightarrow \infty$$

for all $a \in \mathcal{A}$, where $E_{\mathbf{1}_\mathcal{A} \otimes \bigotimes_1^n \mathcal{C}}$ denotes the conditional expectation obtained by evaluating the state ϕ on $\mathcal{A}$.

However the latter condition is well known to be equivalent to the asymptotic completeness of the stationary Markov chain in a scattering theory meaning introduced by Kümmerer and Maassen

in [30]. As mentioned earlier here we followed a variant developed in [20]. The equivalence of the convergence in the norm $\|\cdot\|_2$ above with the property of asymptotic completeness in the scattering theory meaning is stated in Lemma 1.5 of [20]. Hence, proceeding as above, all these meanings of asymptotic completeness coincide. Let us summarize this insight as follows:

Proposition 7.2. The short exact sequence of weak processes associated to a stationary Markov chain is asymptotically complete in the sense of Definition 6.4 if and only if the stationary Markov chain is asymptotically complete in the scattering theory meaning of [30] or [20]. A necessary and sufficient criterion is given by

$$\lim_{n \rightarrow \infty} Z^n(P_{\mathfrak{g}}) = \mathbb{1}_{\mathfrak{h}} \quad (\text{in the strong operator topology})$$

In fact, the last statement is nothing but criterion (5) for asymptotic completeness from Theorem 6.3 applied to this special situation. This reproduces a criterion for asymptotic completeness in terms of the completely positive map Z (which in [20] is called the dual extended transition operator Z' , see [20], Section 4, in particular Theorem 4.3). Because in this case $\mathfrak{g} = \mathbb{C}\Omega_\phi$ is one-dimensional the criterion is also equivalent to the ergodicity of Z , i.e. the fixed point set of Z being equal to $\mathbb{C}\mathbb{1}_{\mathfrak{h}}$, see [20], Section 3, for more details.

The construction of the scattering theory itself with its intertwiners (Møller operators) between a free and a perturbed dynamics requires a two-sided process (i.e. with time variable in $\mathbb{Z}$ for discrete time) as in [30] or the construction of a two-sided extension as in [20] and it would take us too far to include it here. The reader who follows the given sources will recognize that the change of basis studied in Section 6 relating the weak filtration of $(\mathcal{H}, V, \mathfrak{h})$ to the weak filtration of $(\mathcal{G}, V^{\mathcal{G}}, \mathfrak{g})$ and given by the identity on the one-dimensional $\mathfrak{g} = \mathbb{C}\Omega_\phi$ plus the observability map from $\mathfrak{k}$ to $\mathcal{Y}_+$ plus the multi-analytic operator associated to the transfer function (on $\mathcal{U}_+$), is a GNS-version of the Møller operator in the scattering theory, in the same way as the associated weak Markov process is a GNS-version of the stationary Markov chain. If we take into account the identification of $\mathcal{G}$ with the inductive limit of the sequence $(\mathfrak{g} \otimes \bigotimes_1^n \mathcal{P})$ mentioned in Section 2 and note that because $\mathfrak{g} = \mathbb{C}\Omega_\phi$ is one-dimensional we can identify it further with an infinite tensor product $\bigotimes_1^\infty \mathcal{P}$ then we arrive essentially at the version of the scattering theory worked out in [21], Chapter 2.

But the setting of this paper is more general. For example we could also consider stationary Markov chains which are not originally constructed by tensor products of algebras and nevertheless associate weak processes via GNS-construction and apply Proposition 7.1 to find subprocesses (with $\mathfrak{g} = \mathbb{C}\Omega_\phi$). Or we can study higher-dimensional co-invariant subspaces $\mathfrak{g}$. The systematic use of weak processes adds conceptual clarity and cohesion to such investigations.

Acknowledgement: This research has been supported by the EPSRC grant EP/G039275/1.

REFERENCES

- [1] W. Arveson, *Noncommutative dynamics and E-semigroups*. Springer Monographs in Mathematics. Springer, 2003.
- [2] J.A. Ball, V. Bolotnikov, Q. Fang, *Multivariable backward-shift-invariant subspaces and observability operators*. Multidimens. Syst. Signal Process. 18 (2007), no. 4, 191-248.
- [3] J.A. Ball, G. Groenewald and T. Malakorn, *Structured noncommutative multidimensional linear systems*, SIAM J. Control Optim. 44 (2005), no. 4, 1474-1528.
- [4] J.A. Ball, G. Groenewald and T. Malakorn, *Conservative structured noncommutative multidimensional linear systems*. The State Space Method: Generalizations and Applications (Ed. D. Alpay and I. Gohberg), OT 161, Birkhäuser (2006), 179-223.
- [5] J.A. Ball, V. Vinnikov, *Lax-Phillips Scattering and Conservative Linear Systems: A Cuntz-Algebra Multidimensional Setting*. Memoirs of the AMS, vol. 178 (2005), no. 837.
- [6] V. P. Belavkin, *Theory of the control of observable quantum systems*. Autom. Rem. Control, 44 (1983), 178-188.

- [7] V. P. Belavkin, *Nondemolition stochastic calculus in Fock space and nonlinear filtering and control in quantum systems*. Proceedings XXIV Karpacz winter school, R. Guelerak and W. Karwowski, eds., Stochastic methods in mathematics and physics, World Scientific, Singapore (1988), 310-324.
- [8] B.V.R. Bhat, K.R. Parthasarathy, *Kolmogorov's existence theorem for Markov processes in C^* algebras*. K. G. Ramanathan memorial issue. Proc. Indian Acad. Sci. Math. Sci. 104 (1994), no. 1, 253-262.
- [9] B.V.R. Bhat, K.R. Parthasarathy, *Markov dilations of nonconservative dynamical semigroups and a quantum boundary theory*. Ann. Inst. H. Poincaré Probab. Statist. 31 (1995), no. 4, 601-651.
- [10] B.V.R. Bhat, *An index theory for quantum dynamical semigroups*. Trans. Amer. Math. Soc. 348 (1996), no. 2, 561-583.
- [11] B.V.R. Bhat, *Cocycles of CCR flows*. Mem. Amer. Math. Soc. 149 (2001), no. 709.
- [12] L. Bouten, R. van Handel, M. James, *An introduction to quantum filtering*. SIAM J. Control and Optimization 46, no. 6 (2007), 2199-2241.
- [13] L. Bouten, R. van Handel, M. James, *A discrete invitation to quantum filtering and feedback control*. SIAM Review 51, (2009), 239-316.
- [14] O. Bratteli, P. Jorgensen, A. Kishimoto, R. Werner, *Pure States on $\mathcal{O}_d$* . J. Operator Theory 43 (2000), 97-143.
- [15] S. Dey, R. Gohm, *Characteristic functions for ergodic tuples*. Integral Equations and Operator Theory 58 (2007), 43-63.
- [16] S. Dey, R. Gohm, *Characteristic Functions of Liftings*. Journal of Operator Theory 65 (2011), 17-45.
- [17] S. Dey, K. Haria, *Generalized repeated interaction model and transfer functions*, preprint arXiv:1112.4590.
- [18] C. Foias, A.E. Frazho, *The Commutant Lifting Approach to Interpolation Problems*. Operator Theory, Advances and Applications, vol. 44, Birkhäuser, 1990.
- [19] I. Gohberg, P. Lancaster, L. Rodman, *Invariant Subspaces of Matrices with Applications*. Classics in Applied Mathematics, vol. 51, SIAM, 2006.
- [20] R. Gohm, B. Kümmerner, T. Lang, *Noncommutative symbolic coding*. Ergodic Theory and Dynamical Systems 26 (2006), 1521-1548.
- [21] R. Gohm, *Noncommutative Stationary Processes*. Lecture Notes in Mathematics 1839, Springer, 2004.
- [22] R. Gohm, *Kümmerner-Maassen scattering theory and entanglement*. Infinite Dimensional Analysis, Quantum Probability and Related Topics, vol.7(2), World Scientific (2004), 271-280.
- [23] R. Gohm, *Decompositions of Beurling type for E_0 -semigroups*. Banach Center Publications, Vol. 73, Institute of Mathematics, Polish Academy of Sciences, Warszawa (2006), 167-176.
- [24] R. Gohm, *Noncommutative Markov Chains and Multi-Analytic Operators*. J. Math. Anal. Appl., vol. 364(1) (2009), 275-288.
- [25] R. Gohm, *Transfer Functions for Pairs of Wandering Subspaces*. preprint arXiv:1102.5519, to appear in: IWOTA 2010 Proceedings, Operator Theory: Advances and Applications, vol. 221 (2012), 389-402.
- [26] J. Gough, R. Gohm, M. Yanagisawa, *Linear quantum feedback networks*. Phys. Rev. A 78, 062104 (2008)
- [27] J. Gough, M.R. James, *Quantum Feedback Networks: Hamiltonian Formulation*. Commun. Math. Phys., Volume 287 (2009), no. 3, 1109-1132.
- [28] J. Gough, A. Sobolev, *Stochastic Schrödinger equations as Limit of Discrete Filtering*. Open Systems & Information Dyn. 11 (2004), 1-21.
- [29] J.W. Helton, *Discrete time systems, operator models and scattering theory*. J. Functional Analysis 16 (1974), 15-38.
- [30] B. Kümmerner, H. Maassen, *A Scattering Theory for Markov Chains*. Inf. Dim. Analysis, Quantum Prob. and Related Topics, vol.3 (2000), 161-176.
- [31] B. Kümmerner, *Quantum Markov Processes*. A. Buchleitner, K. Hornberger (Eds.), Coherent evolution in noisy environments, Springer LNP 611 (2002), 139-198.

- [32] M. Laca, *Endomorphisms of $\mathcal{B}(\mathcal{H})$ and Cuntz Algebras*. J. Operator Theory 30, no.1 (1993), 85-108.
- [33] G. Popescu, *Isometric Dilations for Infinite Sequences of Noncommuting Operators*. Trans. Amer. Math. Soc. 316 (1989), 523-536.
- [34] G. Popescu, *Characteristic Functions for Infinite Sequences of Noncommuting Operators*. J. Operator Theory 22 (1989), 51-71.
- [35] G. Popescu, *On intertwining dilations for sequences of noncommuting operators*. J. Math. Anal. Appl. 167 (1992), no. 2, 382-402.
- [36] G. Popescu, *Multi-Analytic Operators on Fock Spaces*. Math. Ann. 303 (1995), no. 1, 31-46.

INSTITUTE OF MATHEMATICS AND PHYSICS, ABERYSTWYTH UNIVERSITY, ABERYSTWYTH, SY23 3BZ, UK
E-mail address: `rog@aber.ac.uk`